

Product Data Sheet

Anti-Dog IgE (epsilon chain specific) Antibodies and conjugates

Cat# 30390	Goat Anti-human IgE, unlabeled	Size: 0.1 mg
Cat# 30391	Goat Anti-Dog IgE-HRP conjugate	Size: 0.5 ml
Cat# 30393	Goat Anti-Dog IgE-FITC conjugate	Size: 0.5 ml

Description

Goats were immunized with antigen grade **Dog IgE**. Antibodies have been isolated using ammonium sulfate, ion-exchange, and affinity chromatography to remove non-specific antibodies. Specificity of antibody has been tested using IEP, immunodiffusion, and ELISA. The product reacts with Dog IgE with no significant reactivity with Dog IgG, or other serum proteins. The antibody may recognize other species IgE. Purified anti-Dog IgE antibody is supplied as unlabeled or conjugated to Biotin, FITC, AP or HRP.

Form and Storage

Cat# 30390, unlabeled

The antibody is supplied in PBS, pH 7.4, and 0.05% azide in either **lyophilized** (0.1 mg) or **liquid** form (0.1 mg/0.1 ml). Reconstitute powder in PBS to prepare stock solution. Store at -20oC in suitable aliquots. Stability is ~6-12 months. Do not freeze and thaw.

Cat# 30391, HRP-conjugate

Purified antibody was coupled to HRP (RZ>3.0) using periodate method. The molar enzyme to protein (E/P) ratio = 4.0. The antibody is supplied in stabilizing buffer, 0.1% prolcin-300 as preservative in either **lyophilized** or **liquid** form. Reconstitute powder in PBS in 0.5 ml. Store at 4oC in suitable aliquots. Stability is ~6-12 months. Do not freeze and thaw.

Suggested conjugate dilutions are 1:1,000-1:10,000 ELISA, 1:1K-1:5K for western, and 1:200-1:1000 (IHC).

Cat# 30393, FITC-conjugate

Purified anti-Dog IgE antibody was coupled to FITC at F/P ratio ~3:7. The antibody is supplied in PBS, pH 7.4, 0.2% BSA and 0.05% azide in either **lyophilized** or **liquid** form. Reconstitute powder in PBS in 0.5 ml to prepare stock solution. Store at -20oC in suitable aliquots. Stability is ~6-12 months. Do not freeze and thaw.

Suggested conjugate dilutions are 1:200-1:2000 for immunofluorescence.

Absorption Wavelength: 495 nm

Emission Wavelength: 528 nm

Recommended Working Dilution for ELISA

Working dilution for the specific application should be determined by the investigator to obtain the best conditions. Working solution should be prepared immediately before use and diluted solution should be discarded.

Related Material available for ADI

Dog IgE, IgG, IgE, and IgE ELISA kits

Anti-Dog IgE, IgG, IgE, IgE antibodies and conjugates

ELISA kits for the detection of human Antibodies

Western blot recycling kit (Use the same blot/strip to probe with multiple antibodies) **Strip Antibodies in 5 minutes** (no heating or mercaptoethanol)

MSDS: This product is considered non-hazardous as defined by OSHA (CFR 1910.1200. Nov 25, 1983). It can be disposed of in the drain.

30391

150501A