

Product Data Sheet

Anti-Rat IgE (epsilon chain specific) Antibodies and conjugates

Cat# 50519	Goat Anti-rat IgE, unlabeled	Size: 0.1 mg	Form: Liquid	Powder
Cat# 50520	Goat Anti-rat IgE-HRP conjugate	Size: 0.1 ml	Form: Liquid	Powder
Cat# 50530	Goat Anti-rat IgE-FITC conjugate	Size: 0.1 ml	Form: Liquid	Powder
Cat# 50540	Goat Anti-rat IgE-Biotin conjugate	Size: 0.1 ml	Form: Liquid	Powder

Description

Goats were immunized with antigen grade **rat IgE**. Antibodies have been isolated using ammonium sulfate, ion-exchange, and affinity chromatography to remove non-specific antibodies. Specificity of antibody has been tested using IEP, immunodiffusion, and ELISA. The product reacts with rat IgE with no significant reactivity with rat IgG, or other serum proteins. The antibody may recognize other species IgE. Purified anti-rat IgE antibody is supplied as unlabeled or conjugated to Biotin, FITC, AP or HRP.

Form and Storage

Cat# 10159, unlabeled

The antibody is supplied in PBS, pH 7.4, and 0.05% azide in either **lyophilized** (0.1 mg) or **liquid** form (0.1 mg/0.1 ml). Reconstitute powder in PBS in 0.1 ml to prepare 1 mg/ml solution. Store at -20°C in suitable aliquots. Stability is ~6-12 months. Do not freeze and thaw.

Cat# 10160, HRP-conjugate

Purified antibody was coupled to HRP (RZ>3.0) using periodate method. The molar enzyme to protein (E/P) ratio = 4.0. The antibody is supplied in stabilizing buffer, 0.1% proclin-300 as preservative in , either **lyophilized** (0.1 ml) or **liquid** form (0.1 mg/0.1 ml). Reconstitute powder in PBS in 0.5 ml. Store at 4°C in suitable aliquots. Stability is ~6-12 months. Do not freeze and thaw.

Suggested conjugate dilutions are 1:1,000-1:10,000 ELISA, 1:1K-1:5K for western, and 1:200-1:1000 (IHC).

Cat# 10162, FITC-conjugate

Purified anti-rat IgE antibody was coupled to FITC at F/P ratio ~3:7. The antibody is supplied in PBS, pH 7.4, 0.2% BSA and 0.05% azide in either **lyophilized** (0.5 ml) or **liquid** form (0.5 mg/0.5 ml). Reconstitute powder in PBS in 0.5 ml to prepare 1 mg/ml solution. Store at -20°C in suitable aliquots. Stability is ~6-12 months. Do not freeze and thaw.

Suggested conjugate dilutions are 1:200-1:2000 for immunofluorescence.

Absorption Wavelength: 495 nm
Emission Wavelength: 528 nm

Cat# 10163, Biotin-conjugate

Purified anti-rat IgE antibody was coupled to Biotin using ADI Biotinylation kit # 80300 at F/P ratio ~10-20:1. The antibody is supplied in PBS, pH 7.4, 0.2% BSA and 0.05% azide in either **lyophilized** (0.1 mg) or **liquid** form (0.1 mg/0.1 ml). Reconstitute powder in PBS in 0.5 ml to prepare 1 mg/ml solution. Store at -20°C in suitable aliquots. Stability is ~6-12 months. Do not freeze and thaw.

Suggested conjugate dilutions are 1:5,000-1:30,000 ELISA, 1:2K-1:10K for western.

Recommended Working Dilution for ELISA

Working dilution for the specific application should be determined by the investigator to obtain the best conditions. Working solution should be prepared immediately before use and diluted solution should be discarded.

Related Material available for ADI

Rat IgE, IgG, IgE, and IgE ELISA kits

Anti-rat IgE, IgG, IgE, IgE antibodies and conjugates

ELISA kits for the detection of rat Antibodies

Western blot recycling kit (Use the same blot/strip to probe with multiple antibodies) **Strip Antibodies in 5 minutes** (no heating or mercaptoethanol)

MSDS: This product is considered non-hazardous as defined by OSHA (CFR 1910.1200, Nov 25, 1983). It can be disposed of in the drain.

50519, 50520, 50530, 50540

70226A