

Product Data Sheet

**Anti-Chicken IgG (Fc)-Antibodies and conjugates**

<b>Cat#</b> 60419	Goat Anti-Chicken IgG (Fc), unlabeled	<b>Size:</b> 1 mg	<b>Form:</b> Liquid	Powder
<b>Cat#</b> 60420	Goat Anti-Chicken IgG (Fc)-HRP conjugate	<b>Size:</b> 1 ml	<b>Form:</b> Liquid	Powder
<b>Cat#</b> 60430	Goat Anti-Chicken IgG (Fc)-FITC labeled	<b>Size:</b> 0.5 ml	<b>Form:</b> Liquid	Powder
<b>Cat#</b> 60440	Goat Anti-Chicken IgG (Fc)-Biotin labeled	<b>Size:</b> 0.5 ml	<b>Form:</b> Liquid	Powder

**Description**

Goats were immunized with antigen grade **Chicken IgG (Fc)**. Antibodies have been isolated using ammonium sulfate, ion-exchange, and affinity chromatography to remove non-specific antibodies. Purified anti-Chicken IgG (Fc) is supplied as unlabeled IgG, HRP, FITC and Biotin-conjugate.

Specificity of antibody has been tested using IEP, immunodiffusion, and ELISA. The product reacts with Chicken IgG (Fc) with no significant reactivity with Chicken IgG (Fab2), IgM or other serum proteins. The antibody may recognize other species IgG (Fc).

**Form and Storage**

**Cat# 60419, unlabeled**

The antibody is supplied in PBS, pH 7.4, and 0.05% azide in either **lyophilized** (1 mg) or **liquid** form (1.0 mg/ml). Reconstitute powder in PBS in 1 ml to prepare 1 mg/ml solution. Store at -20oC in suitable aliquots. Stability is ~6-12 months. Do not freeze and thaw.

**Cat# 60420, HRP-conjugate**

Purified antibody was coupled to HRP (RZ>3.0) using periodate method. The molar enzyme to protein (E/P) ratio = 4.0. The antibody is supplied in stabilizing buffer, 0.1% prolcin-300 as preservative in either **lyophilized** (1 ml) or **liquid** form (1 ml). Reconstitute powder in PBS in 1 ml. Store at 4oC in suitable aliquots. Stability is ~6-12 months. Do not freeze and thaw.

Suggested conjugate dilutions are 1:1,000-1:10,000 ELISA, 1:1K-1:5K for western, and 1:200-1:1000 (IHC).

**Cat# 60430, FITC-conjugate**

Purified anti-Chicken IgG (Fc) antibody was coupled to FITC at F/P ratio ~4-5:1. The antibody is supplied in PBS, pH 7.4, 0.2% BSA and 0.05% azide in either **lyophilized** (0.5 ml) or **liquid** form (0.5 mg/0.5 ml). Reconstitute powder in PBS in 0.5 ml to prepare 1

mg/ml solution. Store at -20oC in suitable aliquots. Stability is ~6-12 months. Do not freeze and thaw.

Suggested conjugate dilutions are 1:200-1:2000 for immunofluorescence.

**Absorption Wavelength:** 495 nm

**Emission Wavelength:** 528 nm

**Cat# 60440, Biotin-conjugate**

Purified anti-Chicken IgG (H+L) antibody was coupled to Biotin using ADI Biotinylation kit # 80300 at F/P ratio ~10-20:1. The antibody is supplied in PBS, pH 7.4, 0.2% BSA and 0.05% azide in either **lyophilized** (0.5 mg) or **liquid** form (0.5 mg/0.5 ml). Reconstitute powder in PBS in 0.5 ml to prepare 1 mg/ml solution. Store at -20oC in suitable aliquots. Stability is ~6-12 months. Do not freeze and thaw.

Suggested conjugate dilutions are 1:5,000-1:30,000 ELISA.

**Recommended Working Dilution for ELISA**

Working dilution for the specific application should be determined by the investigator to obtain the best conditions.

Working solution should be prepared immediately before use and diluted solution should be discarded.

in vitro research use only.

**Related Material available for ADI**

**Chicken IgG (Fc), IgG, IgM, and IgE ELISA kits**

Anti-Chicken IgG (Fc), IgG, IgM, IgE antibodies and conjugates

ELISA kits for the detection of Chicken Antibodies

**Western blot recycling kit** (Use the same blot/strip to probe with multiple antibodies) **Strip Antibodies in 5 minutes** (no heating or mercaptoethanol)

60419, 60420, 60430, 60440

70226A