

Product Data Sheet

Anti-Monkey IgG (H+L) Antibodies and conjugates

Cat# 70010	Goat Anti-Monkey IgG (H+L), unlabeled	Size: 0.5 mg	Form: Liquid; Powder
Cat# 70020	Goat Anti-Monkey IgG (H+L)-HRP conjugate	Size: 0.5 ml	Form: Liquid; Powder
Cat# 70024	Goat Anti-Monkey IgG (H+L)-Alk. Phos. conjugate	Size: 0.5 ml	Form: Liquid; Powder
Cat# 70029	Goat Anti-Monkey IgG (H+L)-Biotin conjugate	Size: 0.5 ml	Form: Liquid; Powder
Cat# 70330	Goat Anti-Monkey IgG (H+L)-FITC conjugate	Size: 0.5 ml	Form: Liquid; Powder

Description

Goats were immunized with antigen grade **Monkey IgG (H+L)**. Antibodies have been isolated using ammonium sulfate, ion-exchange, and affinity chromatography to remove non-specific antibodies. Specificity of antibody has been tested using IEP, immunodiffusion, and ELISA. The product reacts with Monkey IgG, IgA and IgM with no significant reactivity with other serum proteins. The antibody may recognize other species IgG (H+L or IgA, IgG, IgM etc). Purified anti-Monkey IgG (H+L) antibody is supplied as unlabeled or conjugated to Biotin, FITC, AP or HRP.

Form and Storage

Cat# 70010, Anti-Monkey IgG (H+L), unlabeled

The antibody is supplied in PBS, pH 7.4, and 0.05% azide in either **lyophilized** (1.0 mg) or **liquid** form (1 mg/ml). Reconstitute powder in PBS in 1.0 ml to prepare 1 mg/ml solution. Store at -20oC in suitable aliquots. Stability is ~6-12 months. Do not freeze and thaw.

Cat# 70020, Anti-Monkey IgG (H+L)-HRP-conjugate

Purified antibody was coupled to HRP (RZ>3.0) using periodate method. The molar enzyme to protein (E/P) ratio = 4.0. The antibody is supplied in stabilizing buffer, 0.1% prolcin-300 as preservative in either **lyophilized** (0.5 ml) or **liquid** form (0.5 ml; 0.5-1 mg/ml). Reconstitute powder in PBS in 1 ml. Store at 4oC in suitable aliquots. Stability is ~6-12 months. Do not freeze and thaw.

Suggested conjugate dilutions are 1:1,000-1:10,000 ELISA, 1:1K-1:5K for western, and 1:200-1:1000 (IHC).

Cat# 10124-AP, AP-conjugate

The conjugate is provided at ~0.5-1 mg/ml as liquid in a stabilizing buffer (50 mM Tris-150 mM NaCl-1 mM MgCl₂, pH 7.5, containing 1% bovine serum albumin, 0.05% sodium azide and 50% glycerol). The product should be **stored at 4°C** and is stable for a minimum of 1 year. Do not store diluted solutions.

Cat# 70029, Anti-Monkey IgG (H+L)-Biotin-conjugate

Purified antibody was coupled to Biotin using ADI Biotinylation kit # 80300 at F/P ratio ~10-20:1. The antibody is supplied in PBS, pH 7.4, 0.2% BSA and 0.05% azide in either **lyophilized** (0.5 mg) or **liquid** form (0.5 mg/0.5 ml). Reconstitute powder in PBS in 0.5 ml to prepare 1 mg/ml solution. Store at -20oC in suitable aliquots. Stability is ~6-12 months. Do not freeze and thaw.

Suggested conjugate dilutions are 1:5,000-1:30,000 ELISA, 1:2K-1:10K for western.

Cat# 70330, FITC-conjugate

Purified antibody was coupled to FITC at F/P ratio ~3:7. The antibody is supplied in PBS, pH 7.4, 0.2% BSA and 0.05% azide in either **lyophilized** (0.5 ml) or **liquid** form (0.5 mg/0.5 ml). Reconstitute powder in PBS in 0.5 ml to prepare 1 mg/ml solution. Store at -20oC in suitable aliquots. Stability is ~6-12 months. Do not freeze and thaw.

Suggested conjugate dilutions are 1:200-1:2000 for immunofluorescence.

Absorption Wavelength: 495 nm

Emission Wavelength: 528 nm

Recommended Working Dilution for ELISA

Working dilution for the specific application should be determined by the investigator to obtain the best conditions. Working solution should be prepared immediately before use and diluted solution should be discarded.

Related Material available for ADI

Monkey IgG, IgM ELISA kits

Anti-Monkey IgG (H+L), IgG, IgG (H+L), IgG (H+L) antibodies and conjugates

ELISA kits for the detection of Monkey Antibodies

Western blot recycling kit (Use the same blot/strip to probe with multiple antibodies) **Strip Antibodies in 5 minutes** (no heating or mercaptoethanol)

70010, 70020, 70024, 70029, 70330

70724A