

Product Specification Sheet

**Adenovirus Antibodies**

<input type="checkbox"/> <b>Cat. ADV12-M</b>	Monoclonal Anti-Adenovirus (many istoypes) hexon IgG	<b>SIZE:</b> 100 ul
<input type="checkbox"/> <b>Cat. ADV12-FITC</b>	Monoclonal Anti-Adenovirus (many istoypes) hexon IgG-FITC conjugate	<b>SIZE:</b> 100 ul

The adenovirus is an ubiquitous pathogen of humans and animals. Adenoviruses are characterized by location inside the cell nucleus, common complement-fixing antigens and marked stability to environmental effects. Adenoviruses are endemic in all populations throughout the year. The infection is spread both through the aerial-droplet route and the routes characteristic for intestinal infections. The incubation period is between five and seven days. Adenoviruses mainly infest respiratory and intestinal mucosa, but also the cornea. They are accumulated in the epithelial cells and regional lymph nodes. Adenoviruses cause the widest variety of illnesses of the known respiratory viruses. The adenovirus infection is the most frequently caused viral disease of the respiratory tract among preschool children (types 1- 5 and 7). Acute diseases of the upper respiratory tract occur predominantly. Pneumonia is the most severe form of adenoviral infection occurring mostly in infants below the age of one. Adenoviruses also cause outbreaks of swimming-pool associated pharyngo conjunctival fever in the summer and epidemics of kerato-conjunctivitis of both children and adults. The intestinal form of adenoviral infection occurs mostly in children below the age of one. An acute adenoviral infection can be detected by virus isolation and/or serology. The serologic tests are particularly important because they document actual infection in the patient and can be applied to large scale epidemiologic investigations. The CF and ELISA tests measure predominantly the antibodies directed against the group-specific determinants on the hexon component.

The hexon is the major capsid protein of the adenovirion and is composed of three identical large polypeptide chains, each containing approximately 1000 residues. The viral shell has a total of 240 such capsomers which assemble in groups of nine to constitute the major part of the 20 triangular surfaces of the adenovirus icosahedron. The major structural features are shared by adenovirus of all serotypes, but the hexon polypeptide, as well as other capsid proteins, differ in size and immunological properties between serotypes. Different forms of hexon from one serotype have also been described.

**Source of Antigen and Antibodies**

<b>Antigen</b>	Adenovirus type 3 (ATCC strain VR847)
<b>Antibody host/type</b>	Mouse, monoclonal IgG1 (Cat # <b>ADV12-M</b> ) unlabeled or Coupled to FITC (# <b>ADV12-FITC</b> )
<b>Secondary Ab</b>	<b>Goat Anti-mouse IgG-HRP conjugate</b> Cat # 40320 (AP, biotin, FITC conjugates also available)
<b>Negative Control Ab</b>	Cat # 20008-1, Mouse (non-immune) Serum IgG, purified, suitable for ELISA, Western, IHC as -ve control

**Isotype Controls**

20102-101	Mouse IgG1 isotype control, purified
20102-101-1	Mouse IgG1 isotype control, purified
20102-101-APC	Mouse IgG1-APC conjugate (isotype control)
20102-101-B	Mouse IgG1-Biotin conjugate (isotype control)
20102-101-F	Mouse IgG1-FITC conjugate (isotype control)
20102-101-FP	Mouse IgG1-FITC-PE conjugate (isotype control)
20102-101-HP	Mouse IgG1-HRP conjugate (isotype control)
20102-101-PC5	Mouse IgG1-PE-Cy5 conjugate (isotype control)
20102-101-PE	Mouse IgG1-PE conjugate (isotype control)

**Form & Storage of Antibodies/Peptide Control**

**Affinity pure IgG and FITC Conjugates**

100 ul       solution       lyophilized powder  
Supplied in **Buffer:** PBS+0.05% azide  
Dissolve powder in 100 ul water.

**Cat# ADV12-FITC, FITC-conjugate**

Purified antibody was coupled to FITC at F/P ratio ~3:7. The antibody is supplied in PBS, pH 7.4, 0.2% BSA and 0.05% azide in either **lyophilized** (0.1 ml) or **liquid** form (0.1 ml). Reconstitute powder in water (0.1 ml to prepare stock solution). Store at -20oC in suitable aliquots. Stability is ~6-12 months. Do not freeze and thaw.

Suggested conjugate dilutions are 1:200-1:2000 for immunofluorescence.

**Absorption** Wavelength: 495 nm  
**Emission** Wavelength: 528 nm

**Stability:** 6-12 months at -20oC or below.  
**Shipping:** 4oC for solutions and room temp for powder

**Recommended Usage**

**Western Blotting** (1:250-1:1000) using Chemiluminescence technique.

**ELISA** (1:10K-1:100K; using 50-100 ng of control peptide/well).

**Histochemistry & Immunofluorescence.** Not tested..

**Specificity & Cross-reactivity**

Adenovirus hexon proteins of at least 21 serotypes including types 40 and 41.

**General References:** Philipson L (1979) Adv. Virus. Res. 25, 357-405; Pettersson U (1971) virology 84, 123-136; Bulanger P (1978) Virology 84, 456-468; Harrison SC (2010) Science 329, 1026-1027;  
\*This product is for in vitro research use only.

**Related material available from ADI**

950-100-AHA	Human Anti-Adenovirus IgA ELISA kit
950-110-AHG	Human Anti-Adenovirus IgG ELISA kit
950-120-AHM	Human Anti-Adenovirus IgM ELISA kit
950-130-AMG	Mouse Anti-Adenovirus IgG ELISA kit
950-140-AMM	Mouse Anti-Adenovirus IgM ELISA kit
ADV11-BTN	Anti-Adenovirus type 2, hexon IgG-Biotin
ADV11-FITC	Anti-Adenovirus type 2, hexon IgG-FITC conjugate
ADV11-HRP	Anti-Adenovirus type 2, hexon IgG-HRP conjugate
ADV11-S	Anti-Adenovirus type 2, hexon antiserum (reacts with 1-7a, 8, 31, 40-41)
ADV12-FITC	Monoclonal Anti-Adenovirus (many isotypes) IgG
ADV12-M	Monoclonal Anti-Adenovirus (many istoypes)
ADV13-M	Monoclonal Anti-Adenovirus type 40 IgG, aff pure
ADV14-M	Monoclonal Anti-Adenovirus type 41 IgG, aff pure
ADV15-M	Monoclonal Anti-Adenovirus type 40/41 IgG,
ADV16-M	Monoclonal Anti-Adenovirus hexon (types 1, 5, 8, 27) IgG
ADV17-M	Monoclonal Anti-Adenovirus type (pan, reacts with all human serotypes) IgG, aff pure
ADV12-M	140521A

Alpha Diagnostic Intl Inc., 6203 Woodlake Center Dr, San Antonio, TX 78244, USA; **Email:** [service@4adi.com](mailto:service@4adi.com)

**India Contact:**

**Life Technologies (India) Pvt. Ltd.**

306, Aggarwal City Mall, Opposite M2K Pitampura, Delhi - 110034 (INDIA). Ph: +91-11-42208000, 42208111, 42208222, Mobile: +91-9810521400, Fax: +91-11-42208444  
Email: [customerservice@lifetechindia.com](mailto:customerservice@lifetechindia.com) Website: [www.lifetechindia.com](http://www.lifetechindia.com)