



Via Sant'anna 131/135
61030 Cartoceto PU (IT)
Telephone + 39 (0) 721830605
FAX +39 (0)721837154
e-mail info@diatheva.com
www.diatheva.com

India Contact:**Life Technologies (India) Pvt. Ltd.**

306, Aggarwal City Mall, Opposite M2K Pitampura,
Delhi – 110034 (INDIA).

Ph: +91-11-42208000, 42208111, 42208222

Mobile: +91-9810521400

Fax: +91-11-42208444

Email: customerservice@lifetechindia.com

Web: www.lifetechindia.com

Rabbit Anti Streptococcus mutans-FITC ANT0088

200µl

Description

Streptococcus mutans is a Gram-positive bacterium that lives in the mouth. The bacterium metabolizes different kinds of carbohydrates, creating an acidic environment in the mouth as a result of this process. This acidic environment in the mouth is what causes the tooth decay. *S. mutans* is considered to be the most cariogenic of all of the oral Streptococci [1].

S. mutans is very important to study, not only because virtually everyone in the world carries it, but also because it has various symptoms that affect our daily lives. As the bacteria develop in the mouth, they cause tooth destruction, impaired speech, difficulty chewing, multiple infections, psychological problems such as low self-esteem, poor social interactions, concentration problems, etc. Though not fatal, tooth decay is one of the most common infectious diseases in humans. Also, cavities caused by the bacteria are the reason for half of all dental visits in the U.S. [2].

The bacteria are most concentrated in the crevices, pits, and fissures that are a normal part of the teeth and surrounding structures. Adults may have a high concentration of *S. mutans* in their mouths. In contrast, infants and children have a smaller concentration, but they are more vulnerable to the bacteria [3]. One of the important virulence properties of these organisms is their ability to form biofilms known as dental plaque on tooth surfaces. Biofilms are sessile bacterial communities adherent to a surface, and their formation occurs in response to a variety of environmental cues. *S. mutans* undergoes a developmental program in response to environmental signals that leads to the expression of new phenotypes that distinguish these sessile cells from planktonic cells (4). Microbiological studies have been conducted in particular in patients undergoing treatment for cancer. Radiotherapy-caused hyposalivation may affect the oral microbiota. Variations in quantity, complexity, and quality of the oral microbiota also occur during chemotherapy, leading to a major imbalance of the ecosystem (5).

Product type	Polyclonal antibody conjugated with FITC
Immunogen	<i>S. mutans</i> ATCC 25175 1010 cells inactivated in glutaraldehyde 2.5%v/v
Source	Rabbit
Reacts with	<i>Streptococcus oralis</i>
Specificity	<i>Streptococcus oralis</i> ; not cross-react against <i>S. mutans</i> ATCC 25175; <i>P. gingivalis</i> ATCC 33277; <i>F. nucleatum</i> ATCC 25586 [6].
Tested applications	FACS; Confocal laser scanning microscopy;
Recommended dilutions	Recommended starting dilutions can vary lot-to-lot. Consult the product information label in the package for lot specific values Note: When using any primary antibody or fluorescence-labelled secondary antibody for the first time, titrate out the antibody to determine which dilution allows the strongest specific signal with the lowest background for your sample [6].
Purity	Polyclonal immunoglobulins purified by protein A affinity chromatography.
Form	Liquid. Supplied in PBS. Neutral pH.
Storage	Shipped at +4°C. When stored at +4°C, the antibody is stable for 18 months. For extended storage, the solution may be frozen at -20°C in working aliquots. Note: Avoid repeated freezing and thawing cycles. Avoid exposure to light.

Reference

- [1] Byers, H.L., E. Tarelli, K. A. Homer, and D. Beighton. "Isolation and characterisation of sialidase from a strain of *Streptococcus oralis*." *Joint Microbial Research Unit*. 49. (2000): 235-244. Web. 16 Dec. 2013.
- [2] Whalan, Rachel H., Simon G. P. Funnell, et al. "Distribution and Genetic Diversity of the ABC Transporter Lipoproteins PiuA and PiaA within *Streptococcus pneumoniae* and Related *Streptococci*." *Journal of Microbiology*. 188.3 (2006): 1031-1038. Web. 16 Dec. 2013.
- [3] Maeda, K., H. Nagata, M. Kuboniwa, et al. "Identification and Characterization of *Porphyromonas gingivalis* Client Proteins That Bind to *Streptococcus oralis* Glyceraldehyde-3-Phosphate Dehydrogenase." *Infection and Immunity*. 81.3 (2013): 753-763. Web. 16 Dec. 2013.
- [4] Wilder, Jennifer, Michelle Ramanathan, et al. "*Streptococcus oralis* meningitis following spinal anaesthesia." *Association of Anesthetist of Great Britain and Ireland*. 0092. (2013): n. page. Web. 16 Dec. 2013.
- [5] "How EvoraPro Oral Probiotics Aid Oral Health." *Evroa Oral Probiotics*. N.p., n. d. Web. 16 Dec. 2013.
- [6] Manti A, Ciandrini E, Campana R, Dominici S, Ciacci C, Federici S, Sisti D, Rocchi MB, Papa S, Baffone W. **A dual-species microbial model for studying the dynamics between oral streptococci and periodontal pathogens during biofilm development on titanium surfaces by flow cytometry.** *Res Microbiol.* 2016 Jun;167(5):393-402. doi: 10.1016/j.resmic.2016.03.004. Epub 2016 Mar 28