

Product Specification Sheet

Human Apolipoprotein E3 (ApoE3) Antibodies

Cat. # APOE31-S	Rabbit Anti-Human ApoE3 antiserum # 1	SIZE: 100 ul
Cat. # APOE31-C	Human ApoE3 protein Western Blot +ve control	SIZE: 100 ul

Apolipoprotein E (apo E), a component of lipoproteins produced by the liver and in circulating macrophages, plays a critical role in the reverse transport of cholesterol to the liver via the circulation. ApoE is also expressed in the brain and in response to injury in both the peripheral and central nervous systems. **ApoE** is a single polypeptide chain of 299 amino acids (~34 kDa) consisting of two independently folded functional domains. The C-terminal domain contains the major lipid-binding region. The N-terminal domain exists in the lipid-free state as a four-helix bundle of amphipathic α -helices and contains the LDLR-binding region (amino acids 136–150 in helix), which coincides with a heparin-binding site. Humans and mice lacking apoE cannot clear remnant lipoproteins from the plasma and are at increased risk for atherosclerosis. ApoE is a high-affinity ligand for the low density lipoprotein receptor (LDLR) family and for cell-surface heparan sulfate proteoglycans. Defective binding of apoE to receptors causes cholesterol-rich lipoprotein particles to accumulate in the plasma and is the mechanism of type III hyperlipoproteinemia, a genetic disorder characterized by elevated plasma cholesterol and triglyceride levels and accelerated coronary artery disease. In humans, apoE has three major isoforms: **Apo E2** (Cys112, Cys158), **Apo E3** (Cys112, Arg158), **Apo E4** (Arg112, Arg158), products of alleles at a single gene locus. Although apoE4 is neither necessary nor sufficient to cause AD, inheritance of apoE4 is a significant risk factor for late onset AD, decreasing the age of onset and the duration of disease. Due primarily to this genetic linkage, the role of apoE in the pathogenesis AD is being actively studied.

Source of Antigen, Antibodies, and positive controls

Antigen	Purified recombinant human ApoE3 (cat # APOE35-R)
Ab Host/type	Rabbit, Polyclonal unpurified antiserum (#APOE31-S)
2-Ab	Cat # 20320, goat anti-rabbit IgG-HRP (AP, biotin, FITC conjugates also available).
-ve control IgG	# 20009-1, Rabbit (non-immune) IgG, purified, suitable for ELISA, Western, IHC as -ve control

Human **ApoE3 protein** for Western blot +ve control (Cat # **APOE31-C**) is supplied in SDS-PAGE sample buffer (reduced). Load 10 ul/lane of **APOE31-C** for good visibility with antibody Cat # **APOE31-S**. Store at -20oC in suitable size aliquots. SDS may crystallize in cold conditions. It should redissolve by warming before taking it from the stock. It should be heated once prior to loading on gels. If the product has been stored for several weeks, then it may be preferable to add 5 ul of fresh 2x sample buffer per 10 ul of the **APOE31-C** solution prior to heating and loading on gels. This preparation is not biologically active. It is not suitable for ELISA or other applications where native protein is required. This preparation is intended for qualitative purpose and not to serve as standard of known concentration. Do not freeze, thaw, or heat repeatedly

Form & Storage of Antibodies/Peptide Control

Antiserum (unpurified, undiluted)

100 ul/vial solution lyophilized powder
contains 0.05% sodium azide **Reconstitute powder** 100 ul PBS

Storage

Short-term: unopened, undiluted liquid vials for less than a week at 4oC.

Long-term: at -20C or below in suitable aliquots after reconstitution. Do not freeze and thaw and store working, diluted solutions.

Stability: 6-12 months at -20oC or below.

Shipping: 4oC for solutions and room temp for powder.

Recommended Usage

Western Blotting 1:1K for neat serum using Chemiluminescence technique). Human ApoE3 is approx. ~34 kDa.

ELISA (1:10K-1:100K; using 50-100 ng of control peptide/well).

Histochemistry & Immunofluorescence: not tested. We recommend a dilution of 1:200 to 1:500.

Specificity & Cross-reactivity

The APOE31-S reacts with human ApoE3 isoforms is western and ELISA. It is expected to react with human other ApoE isoforms (E2, and E4). Antibody crossreactivity in various species is not established. **cat # APOE31-C** protein control should be used a positive control.

General References:

1. Alzheimer' Reports (1998)1, 119-128, Neuroscience Lett. (1992) 135, 235-238; Acta Neuropathologica (1996) 92, 378-385; Kelly ME et al (1994) cell Immunol. 159, 124; Mahley RW et al (1988) Science 240, 622; Rall SC et al (1986) Methods Enzymol. 128, 273

*This product is for In vitro research use only.

Related material available from ADI

Ant-Beta amyloid 1-40, 1-42, APP, Parkin, Synucleins (α , β , γ), Presenilins 1, 2, ERAB

APOE31-S-C 71213A

India Contact:

Life Technologies (India) Pvt. Ltd.

306, Aggarwal City Mall, Opposite M2K Pitampura, Delhi – 110034 (INDIA). Ph: +91-11-42208000, 42208111, 42208222, Mobile: +91-9810521400, Fax: +91-11-42208444
Email: customerservice@lifetechindia.com Website: www.lifetechindia.com