

Aquaporin 12 (AQP12) antibodies

| | | |
|-----------------------|--|---------------------|
| Cat # AQP126-P | Human AQP12 control/blocking peptide | SIZE: 100 ug |
| Cat # AQP126-A | Rabbit Anti- Human AQP12 IgG (aff pure) | SIZE: 100 ug |

Water is a critical component of all living cells. Interestingly, tissue membranes show a great degree of water permeability. Mammalian red cells, renal proximal tubules, and descending thin limb of Henle are extraordinarily permeable to water. Water crosses hydrophobic plasma membranes either by simple diffusion or through a facilitative transport mechanism mediated by special protein "aquaporin".
AQP12 (mouse 290 aa; human 295 aa).

SUBCELLULAR LOCATION: Membrane; Multi-pass membrane protein (Potential).

TISSUE SPECIFICITY: Restricted to pancreatic acinar cells.

SIMILARITY: Belongs to the MIP/aquaporin (TC 1.A.8) family. AQP11/AQP12 subfamily.

Protein name Aquaporin-12A , Aqp12

Synonym AQP-12

Gene name Aqp12a

Source of Antigen and Antibodies

| | |
|----------------------------|---|
| Antigen | 18-aa peptide of Human AQP12 (Protein Accession # Q8IXF9; ref 1); designated as AQP126-P control/blocking peptide conjugated to KLH; epitope location ~ C-terminus |
| Antibody host/type | Rabbit, Polyclonal IgG (Cat # AQP126-A), purified over antigen-Agarose |
| 2-Ab | Cat # 20320, goat anti-rabbit IgG-HRP (AP, biotin, FITC conjugates also available). |
| Negative Control Ab | Non-immune rabbit IgG (Cat # 20009-1) to be used as -ve control for ELISA, WB, IHC etc. |

Form & Storage of Antibodies/Peptide Control

Affinity pure IgG

100 ug/100ul solution lyophilized powder
Supplied in **Buffer:** PBS+0.1% BSA
Reconstitute powder in PBS at 1mg/ml

Control/blocking peptide

100 ug/100 ul solution lyophilized powder
Supplied in **Buffer:** PBS pH 7.5,
Reconstitute powder in PBS at 1 mg/ml.

Storage

Short-term: unopened, undiluted liquid vials at -20OC and powder at 4oC or -20oC..

Long-term: at -20C or below in suitable aliquots after reconstitution. Do not freeze and thaw and store working, diluted solutions.

Stability: 6-12 months at -20oC or below.

Shipping: 4oC for solutions and room temp for powder.

Recommended Usage

We recommend the use of 0.5-1% milk in all primary/secondary antibody-enzyme conjugate incubations in order to suppress non-specific bands.

Western Blotting 1:1K-5K for neat serum and 1-10 ug/ml for affinity pure antibody using ECL.

ELISA: Control peptide can be used to coat ELISA plates at 1 ug/ml and detected with antibodies (1:10-50K for neat serum and 0.5-1 ug/ml for affinity pure).

Histochemistry & Immunofluorescence: not tested. we recommend the use of affinity purified IgG at 2-10 ug/ml in paraformaldehyde fixed sections of tissues.

Specificity & Cross-reactivity

Human AQP126-P peptide sequence has no homology to any other known eukaryotic protein sequences. Control peptide, because of its low mol. Wt (<3 kDa), is not suitable for Western. It should be used for ELISA or antibody blocking experiments (use 5-10 ug control peptide per 1 ug of aff pure IgG or 1 ul antiserum) to confirm antibody specificity (see detailed protocol at: www.4adi.com\data/abblock.html).

General References: 1) Itoh T, et al (2005) BBRC 330(4), 832-838.

**This product is for In vitro research use only.*

Related material available from ADI

Antibodies for AQP1-12 rUT2), VMAT1, VMAT2, Vasopressin receptor (AVPR-V1 and V2)

Western blot Recycling kit (probe same blot with multiple antibodies)

AQP126-S-A 70910A