

Product Specification Sheet

Aquaporin 2 (AQP-CD or WCH-CD) Antibodies

Cat # AQP22-P	Rat AQP2 control/blocking peptide	SIZE: 100 ug
Cat # AQP22-S	Rabbit Anti-rat AQP2 antiserum #1	SIZE: 100 ul
Cat # AQP22-A	Rabbit Anti-rat AQP2 IgG #1 (Aff pure)	SIZE: 100 ug

Water is a critical component of all living cells. Interestingly, tissue membranes show a great degree of water permeability. Mammalian red cells, renal proximal tubules, and descending thin limb of Henle are extraordinarily permeable to water. Water crosses hydrophobic plasma membranes either by simple diffusion or through a facilitative transport mechanism mediated by special protein "aquaporin". Over the last decade, genes for several members of aquaporin family (**AQP1-9 and AQPAP**) have been cloned, expressed, and their distribution studied in many tissues. **AQP2 or AQP-CD or WCH-CD** (mouse, rat, and human 271 aa; human gene locus 12q13) is the vasopressin-regulated water channel of the apical membrane of collecting duct cells. AQP families of proteins are predicted to contain six transmembrane domains. The N and C-terminus are predicted to be cytoplasmic. Defects in AQP2 are a cause of an autosomal dominant form of nephrogenic diabetes insipidus (NDI).

Source of Antigen and Antibodies

Antigen	15aa peptide of Rat AQP2 ; Designated (AQP22-P or control peptide /blocking peptide) conjugated to KLH; epitope location ~ C-terminus
Ab Host/type	Rabbit, Polyclonal unpurified antiserum (# AQP22-S) and IgG, purified over antigen-agarose (Cat # AQP22-A)
2-Ab	Cat # 20320, goat anti-rabbit IgG-HRP (AP, biotin, FITC conjugates also available).
-ve control IgG	# 20009-1, Rabbit (non-immune) IgG, purified, suitable for ELISA, Western, IHC as -ve control

Form & Storage of Antibodies/Peptide Control

Antiserum (unpurified, undiluted)

100 ul/vial solution contains 0.05% sodium azide
50 ul/vial lyophilized powder
Reconstitute powder 50 ul or 100 ul PBS

Affinity pure IgG

100 ug/100ul solution
50 ug/50 ul lyophilized powder
Buffer: PBS+0.1% BSA+
Reconstitute powder in PBS at 1mg/ml

Control/blocking peptide

100 ug/100 ul solution
50 ug/50 ul lyophilized powder
Buffer: PBS pH 7.5,
Reconstitute powder in PBS at 1 mg/ml.

Storage

Short-term: unopened, undiluted liquid vials for less than a week at 4oC.

Long-term: at -20C or below in suitable aliquots after reconstitution. Do not freeze and thaw and store working, diluted solutions.

Stability: 6-12 months at -20oC or below.

Shipping: 4oC for solutions and room temp for powder.

Recommended Usage

Western Blotting (1:1K-5K for neat serum and 1-10 ug/ml for affinity pure antibody using ECL). A similar pattern is seen in human kidney. (see published refs using this antibody in 2).

ELISA: Control peptide can be used to coat ELISA plates at 1 ug/ml and detected with antibodies (1:10-50K for neat serum and 0.5-1 ug/ml for affinity pure).

Histochemistry & Immunofluorescence: we recommend the use of affinity purified antibody at 2-10 ug/ml in paraformaldehyde fixed sections of tissues. (see published refs using this antibody in 2).

Specificity & Cross-reactivity

Rat AQP22-P peptide is 100% conserved in mouse, 93% in human and ovine AQP2. Antibody cross-reactivity in various species has not been studied. Control peptide, because of its low mol. Wt (<3 kDa), is not suitable for Western. It should be used for ELISA or antibody blocking experiments (use 5-10 ug control peptide per 1 ug of aff pure IgG or 1 ul antiserum) to confirm antibody specificity.

General References: 1. Fushimi K, (1993) Nature 361, 549; (2) Nielsen, S (1993) PNAS 90, 11663-11667; Deen, PMT (1995) J. Clin. Invest. 95, 2291; (3) Sasaki, S (1994) J. Clin. Invest. 93, 1250.

(2) Citations of ADI's Antibodies (see web site for updated list)

Buemi M, 2004, Am Journal of Hypertension 17, 1170-1178, WB, , Leung JCK, 2005, Nephrology 10, 63-72, WB, , IHC, Ferguson CJ, 2000, Kidney Intl. 64, 1755-1764, WB, , IF Ross, MJ, 2001, J Am Soc Nephrol 2001 12: 2645-2651, , IHC, Saito O, 2003, Am J Physiol Renal Physiol, 284, 1164-1170, , IHC, Schneider A, 2002, Am J Physiol Renal Physiol : 284: 411, IHC, van Vonderen IK, 2004, Domestic Animal Endocrinol. 27, 141-153

*This product is for In vitro research use only.

Related material available from ADI

Antibodies for AQP1-9 & AQP-AP, & rUT2; VMAT1/2

AQP22-S-A-P 80709A

India Contact:

Life Technologies (India) Pvt. Ltd.

306, Aggarwal City Mall, Opposite M2K Pitampura, Delhi - 110034 (INDIA). Ph: +91-11-42208000, 42208111, 42208222, Mobile: +91-9810521400, Fax: +91-11-42208444
Email: customerservice@lifetechindia.com Website: www.lifetechindia.com