

Product Specification Sheet

Arginine Vasopressin Receptor (AVP-V2) Antibodies

Cat # AVPV21-P	Rat AVP-V2 Control Peptide	SIZE: 100 ug
Cat # AVPV21-S	Rabbit Anti-Rat AVP-V2 antiserum	SIZE: 100 ul
Cat # AVPV21-A	Rabbit Anti-Rat AVP-V2 Ig G, Aff pure	SIZE: 100 ug

Vasopressin (**AVP**, Arginine-8-Vasopressin), the antidiuretic hormone is cyclic nonapeptide involved in the homeostasis of body fluid osmolality, blood volume, vascular tone, and blood pressure. Specific actions of AVP include inhibition of diuresis, contraction of smooth muscle, stimulation of liver glycogenesis, and modulation of ACTH release from pituitary. AVP belongs to the family of vasoactive and mitogenic peptide involved in normal and pathological cell growth and differentiation.

AVP exerts its action through binding to specific membrane receptors coupled to distinct second messengers. There are 3 types of AVP receptors: **V1a, V1b, and V2** subtypes. The V2 receptor stimulates adenylyl cyclase and protein Kinase A, V1 activate phospholipase A2.C, and D, resulting into production of IP3 and DAG, the mobilization of intracellular calcium, the influx of extracellular calcium, the activation of protein Kinase C, and protein phosphorylation. The V1a receptors mediate vasoconstriction and hepatic gluconeogenesis platelet aggregation, coagulation factor release. V1a receptors are found in vascular smooth muscle, hepatocytes, blood platelets, lymphocytes and monocyte, type II pneumocytes, adrenal cortex, brain, reproductive organs, retinal epithelium, renal mesangial cells.

AVP receptors are members of the G-protein coupled receptors with putative 7 transmembrane domains. The sizes of various AVP receptors are V1a (rat, 424 AA; human, 418 AA); V1b (rat, 421 AA; human, 424 AA); V2 (rat, 371 AA; human 371 AA). The N-terminus and C-terminus are predicted to be extracellular and cytoplasmic, respectively.

Source of Antigen and Antibodies

Antigen	21aa peptide of rat AVPV2 receptor ; Designated (AVPV21-P or control peptide). Epitope location ~ C-terminus, Cytoplasmic
Ab Host/type	Rabbit, polyclonal Unpurified antiserum (cat # AVPV21-S) Aff pure IgG (cat #AVPV21-A)purified over the antigen column
2ab	Cat # 20320, goat anti-rabbit IgG-HRP (AP, biotin, FITC conjugates also available
-ve control	# 20009-1, Rabbit (non-immune) IgG, purified, suitable for ELISA, Western, IHC as -ve control

Form & Storage of Antibodies/Peptide Contro

Antiserum (unpurified)
100ul solution lyophilized powder
Supplied in Buffer: 0.05% azide
Reconstitute powder in 100 ul PBS

Affinity pure IgG
100 ug/100ul solution lyophilized powder
Supplied in **Buffer:** PBS+0.1% BSA
Reconstitute powder in PBS at 1mg/ml

Control/blocking peptide

100 ug/100 ul solution lyophilized powder
Supplied in Buffer: PBS pH 7.5,

Reconstitute powder in PBS at 1 mg/ml.

Storage

Short-term: unopened, undiluted liquid vials at 20°C and powder at 4°C or -20°C..

Long-term: at -20°C or below in suitable aliquots after reconstitution. Do not freeze and thaw and store working, diluted solutions.

Stability: 6-12 months at -20°C or below.

Shipping: 4°C for solutions and room temp for powder

Recommended Usage

Western Blotting (1:1K-5K for neat serum and 1-10 ug/ml for affinity pure antibody using Chemiluminescence technique).

ELISA: Control peptide can be used to coat ELISA plates at 1 ug/ml and detected with antibodies (1:10-50K for neat serum and 0.5-1 ug/ml for affinity pure).

Histochemistry & Immunofluorescence: Not tested. we recommend the use of affinity purified antibody at 2-10 ug/ml in paraformaldehyde fixed sections of tissues. Neat serum can be used at 1:500 or more.

Specificity & Cross-reactivity

The 21 aa rat AVPV21-P has homology with mouse (95%), human (90%), pig (80%), and bovine (76%) V2 receptors. It has no significant similarity with AVP V1a or V1b receptors. Antibody crossreactivity in various species is not established. Control peptide, because of its low mol. Wt (<3 kDa), is not suitable for Western. It should be used for ELISA or antibody blocking experiments (use 5-10 ug control peptide per 1 ug of aff pure IgG or 1 ul antiserum) to confirm antibody specificity (see detailed protocol at:the web site).

General References: 1. Lolait, S et al (1992) Nature 357, 336-339; Firsov D et al (1994) Pflugers Arch 429, 79-89; 2. Nonoguchi, H et al (1995) J. Clin. Invest. 96, 1768-1778.

Citations of ADI's antibodies for Arginine Vasopressin Receptors (see updated list at the web site)

*This product is for In vitro research use only.

Related material available from ADI

Antibodies for AVP-V2 and AVP-V1a (please call for an update)

Antibodies to AQP1-5, and Urea Transporter

VMAT1, VMAT2

AVPV1A12-S-A-P

71215S

India Contact:

Life Technologies (India) Pvt. Ltd.

306, Aggarwal City Mall, Opposite M2K Pitampura, Delhi – 110034 (INDIA). Ph: +91-11-42208000, 42208111, 42208222, Mobile: +91-9810521400, Fax: +91-11-42208444
Email: customerservice@lifetechindia.com Website: www.lifetechindia.com