

Product Data Sheet

- Cat # CFP101-M Monoclonal Anti-M. tuberculosis 10 Kda cultural filtrate protein (CFP10) IgG #1 **Size:** 100 ul
 Cat # CFP102-M Monoclonal Anti-M. tuberculosis 10 Kda cultural filtrate protein (CFP10) IgG #2 **Size:** 100 ul

Tuberculosis, MTB, or TB (short for tubercle bacillus) is a common, and in many cases lethal, infectious disease caused by various strains of mycobacteria, usually *Mycobacterium tuberculosis*. Tuberculosis typically attacks the lungs, but can also affect other parts of the body. Individuals with HIV are at risk for infection by tuberculosis due to their impaired immune system. The only currently available vaccine as of 2012 is bacillus Calmette–Guérin (**BCG** with live attenuated bacteria) which, while it is effective against disseminated disease in childhood, confers inconsistent protection against contracting pulmonary TB. A number of new TB vaccines are currently in phase I and II clinical trials. MVA85A (modified vaccinia Ankara 85A) is a subunit vaccine to BCG.

The closely related proteins of the **antigen 85 complex** are major secreted products of mycobacteria. Three closely related components, termed antigens **85A, 85B, and 85C**, have been demonstrated in *M. bovis* BCG and *M. tuberculosis*. Although the antigens are genetically distinct, they are highly homologous and cross-react with polyclonal and monoclonal antibodies raised against individual components. Sequence analysis revealed 85% identity between the *M. bovis* BCG 85A and 85B. Many other mycobacterial antigens have been identified, such as 71, 65, 38, 23, 19, 16, 14 and 12-kDa proteins. The **38-kDa** protein is an immunodominant lipoprotein antigen is specific only for the *M. tuberculosis* complex. It is the most extensively studied antigen. The 16-kDa antigen is an immunodominant antigen, frequently called 14 kDa, related to the family of low molecular weight heat-shock proteins. This antigen contains B-cell epitopes specific for the *M. tuberculosis* complex.

Mycobacterium antigen **ESAT-6** has been isolated from low molecular weight fractions of the **shot-term-culture filtrate (ST-CF)** and it can easily be detected in tuberculosis patients. This antigen includes many epitopes detectable in the serum of most patients with tuberculosis (more than 90%). Recently it was shown that ESAT-6 is very potential as diagnostic for differentiation between the mycobacterial infection and BCG vaccination.

CFP-10 also known as ESAT-6-like protein esxB or secreted antigenic protein MTSA-10 or 10 kDa culture filtrate antigen CFP-10 is a protein that is encoded by the esxB gene. CFP-10 is a 10 kDa secreted antigen from *Mycobacterium tuberculosis*. It forms a 1:1 heterodimeric complex with ESAT-6. Both genes are expressed from the RD1 region of the bacterial genome and play a key role in the virulence of the infection. **CFP10 (71-85 aa; EISTNIRQAGVQYSR)** is a 15-mer peptide is fragment of (CFP)10 protein, which is secreted from mycobacterium tuberculosis 10-kDa culture filtrate stimulate IFN- γ production and CTL activity by CD4+ and CD8+ cells, from persons expressing a spectrum of MHC molecules. This peptide is an excellent candidate for inclusion in a subunit tuberculosis vaccine.

| | |
|------------------------|---|
| Antigen | Recombinant <i>M. tuberculosis</i> CFP10 protein H37RV |
| Ab Host/type | Mouse, Monoclonal , IgG1, cat # CFP101-M) & #CFP102-M (IgG2b) Supplied in PBS, pH 7.5 containing 0.05% azide, |
| 2-ab | Goat Anti-mouse IgG-HRP conjugate Cat # 40320 (AP, biotin, FITC conjugates also available) |
| -ve control IgG | Cat # 20008-1, Mouse (non-immune) Serum IgG, purified, suitable for ELISA, Western, IHC as –ve control |

Store antibody at 40oC for short term use and frozen at -20oC or below for long term storage. Reconstitute powder in water and store frozen.

Usage

CFP101-M is recommended for ELISA to detect CFP-10 protein. It is optimized for coating and detection with the CFP102-M antibody. Other applications such as western and IHC can be explored as well.

References: Renshaw PS (2005) EMBO J. 24, 2491-2498; Meher Ak (2006) FEBS J. 273, 1445-1462; Sorensen AL (1995) Inf. Immun. 63, 1710-1717; Berthet FX (1998) Microbiol. 144, 3195-3203; Brodin P (2006) Inf. Immun. 74, 88-98; Renshaw PS (2002) JBC 277, 21598-21603

This item is for LABORATORY RESEARCH USE ONLY.

Related Items

| Catalog# | ProdDescription |
|-----------|--|
| HSP651-M | Monoclonal Anti-M. tuberculosis Heat Shock Protein 65 (hsp65/groEL-2/Cpn60-2) IgG |
| HSP651-P | Heat shock protein (M. leprae HSP65; 417-429) specific P62 peptide (LLQAAPALDKLKL, MW:1393.7) |
| HSP652-P | Heat shock protein (M. leprae/M. tuberculosis HSP65; 417-429) P38 peptide (AGGGVTLQAAPALD, MW:1353.5) |
| HSP653-P | Heat shock protein (M. leprae HSP65; 343-355) P61 peptide (RVAQIRTEIENS, MW:1530.7) |
| HSP654-P | Heat shock protein (M. bovis HSP65; 243-255) indicator peptide in HLA-DQ2 binding assays (KPLLLIAEDVEGEY, MW:1588.8) |
| HSP701-M | Monoclonal Anti-M. tuberculosis Heat Shock Protein 70 (hsp70/Dnak/ML2496) IgG |
| MTB061-M | Monoclonal Anti-Mycobacterium tuberculosis antigen (6kDa/ESAT-6) IgG |
| MTB06-R | Recombinant purified (E. coli) Mycobacterium tuberculosis antigen (6kDa/ESAT-6) |
| MTB161-M | Monoclonal Anti-Mycobacterium tuberculosis antigen (16kDa/HspX) IgG |
| MTB16-R | Recombinant purified (E. coli) Mycobacterium tuberculosis antigen (16kDa/HspX) |
| MTB381-M | Monoclonal Anti-Mycobacterium tuberculosis antigen 38kDa/Ag85B IgG |
| MTB38-R | Recombinant purified (E. coli) Mycobacterium tuberculosis antigen (38kDa/Ag85B) |
| MTB6381-S | Anti-M. Tuberculosis antigens (6kDa/ESAT+16kDa+38kDa/Ag85b mixture of proteins antiserum) |
| RP-628 | Recombinant purified Mycobacterium Tuberculosis Heat Shock Protein 70 (hsp70/Dnak/ML2496) |
| RP-999 | Recombinant purified Mycobacterium Tuberculosis major secretory protein Antigen 85B (38kDa Antigen, Ag85b) |

CFP101-MP

Alpha Diagnostic Intl Inc., 6203 Woodlake Center Dr, San Antonio, TX 78244, USA;

India Contact:

Life Technologies (India) Pvt. Ltd.

306, Aggarwal City Mall, Opposite M2K Pitampura, Delhi – 110034 (INDIA). Ph: +91-11-42208000, 42208111, 42208222, Mobile: +91-9810521400

Fax: +91-11-42208444 Email: customerservice@lifetechindia.com Website: www.lifetechindia.com