

Product Specification Sheet

Chloride Channel-2 (CLC-2 or CLCN2) Antibodies

Cat. # CLC21-P	Rat CLC-2 Control Peptide	SIZE: 100 ug
Cat. # CLC21-S	Rabbit Anti-rat CLC-2 antiserum	SIZE: 100 ul
Cat. # CLC21-A	Rabbit Anti-rat CLC-2 Ig G (affinity pure)	SIZE: 100 ug

Chloride is a critical component of all living cells. Voltage-gated chloride channels regulate cellular traffic of chloride ion. The chloride channels (CIC or CLC) performs several functions including the regulation of cell volume, membrane potential stabilization, signal transduction, and transepithelial transport. Mutations in CIC genes have been linked with several human diseases including myotonias (Thomsen's disease), cystic fibrosis, Bartters syndrome type III, Dent's disease, and X-linked recessive nephrolithiasis. In mammals, CLC proteins form a superfamily of at least 9 different genes (CLC1-7 also known as CLCN1-7 and CLK1-2 or CLCKa and CLCKb). Additional forms of these proteins are obtained by alternative splicing. All CLC proteins (~700-1000 aa) are predicted to contain 10 (possibly 12) transmembrane domains. Except CLC-1 and CLC-K1/K2 that are specific for kidney, most other CLC are widely distributed in various tissues.

Rat CLC-2 is 887 aa (or 907 aa) membrane protein (human CLC-2 898 aa) (1). Alternatively spliced short form (491 aa) have also been found. CLC-2 is ubiquitously expressed. Defects in CLC1 (CLCN1) are the cause of autosomal recessive generalized myotonia (Becker's disease) (RGM) and autosomal dominant myotonia congenita (Thomsen's disease; MC) which are characterized by skeletal muscle stiffness (delayed relaxation) that is a result of muscle membrane hyperexcitability

Source of Antigen and Antibodies

Antigen	22aa peptide of rat CL21; Designated (CLC21-P or control peptide) . epitope location ~ C-terminus
Ab Host/type	Rabbit, polyclonal Unpurified antiserum (cat # CLC21-S) Aff pure IgG (cat # CLC21-A) purified over antigen-agarose column
2-ab	Goat Anti-rabbit IgG-HRP cat # 20320 (AP, biotin, FITC conjugates also available)
-ve control IgG	# 20009-1, Rabbit (non-immune) IgG, purified, suitable for ELISA, Western, IHC as -ve control

Form & Storage of Antibodies/Peptide Control

Antiserum (unpurified)
100ul solution lyophilized powder
Supplied 0.05% azide, **Reconstitute** powder in 100 ul PBS

Affinity pure IgG
100 ug/100ul solution lyophilized powder
Supplied in **Buffer:** PBS+0.1% BSA
Reconstitute powder in PBS at 1mg/ml

Control/blocking peptide
100 ug/100 ul solution lyophilized powder
Supplied in **Buffer:** PBS pH 7.5,
Reconstitute powder in PBS at 1 mg/ml.

Storage

Short-term: unopened, undiluted liquid vials at -20OC and powder at 4oC or -20oC..

Long-term: at -20C or below in suitable aliquots after reconstitution. Do not freeze and thaw and store working, diluted solutions.

Stability: 6-12 months at -20oC or below.

Shipping: 4oC for solutions and room temp for powder

Recommended Usage

Western Blotting (1:1K-5K for neat serum and 1-10 ug/ml for affinity pure antibody using ECL technique). An antibody made to the C-terminal region has detected multiple CLC-2 bands between 80-90 kDa bands in various tissues. See refs in 2.

ELISA: Control peptide can be used to coat ELISA plates at 1 ug/ml and detected with antibodies (1:10-50K for neat serum and 0.5-1 ug/ml for affinity pure).

Histochemistry & Immunofluorescence: We recommend the use of affinity purified antibody at 1-20 ug/ml in paraformaldehyde fixed sections of tissues. See refs in 2.

Specificity & Cross-reactivity

The 22 AA rat CLC21-P control peptide is 100% conserved in mouse, 95% in pig, rabbit and 90% in human CLC-2. No significant sequence homology is detected with other CLCs or other proteins. Actual cross-reactivity of antibodies in various species has not been studied. The CLC21-P control peptide is available to confirm specificity of antibodies. Control peptide, because of its low mol. Wt (<3 kDa), is not suitable for Western. It should be used for ELISA or antibody blocking experiments (use 5-10 ug control peptide per 1 ug of aff pure IgG or 1 ul antiserum) to confirm antibody specificity (see detailed protocol at the web site).

General References: Chu S et al (1997) Nucl Acid Res. 25, 4153; Chu S et al (1994) 24, 3453; Cid LP et al (1995) Hum. Mol. Genet. 4, 407; Thiemann A et al (1992) Nature 356, 57-60, Murray CB et al (1995) Am. J. Respir. Cell. Mol. Biol. 12, 597; Murray CB et al (1996) AJP, 271, L829

2. Citations for ADI Antibodies (see updates at the web site)

Hori K 2004 FEBS Lett. 575, 105-108 WB IHC
Gustincich S, 2004, Proc. Natl. Acad. Sci. 101, 5069-5074, IHC

*This product is for In vitro research use only.

Related material available from ADI

Antibodies CLC1-7 and CLC-K1; KCCL1-3; AQP-9 and RUT; OCT and OAT, AE-3, and NACX

CLC21-S-A-P 71217A

Alpha Diagnostic Intl Inc., 6203 Woodlake Center Dr, S an Antonio, T X 7 8 24 4 , U S A;

India Contact:

Life Technologies (India) Pvt. Ltd.

306, Aggarwal City Mall, Opposite M2K Pitampura, Delhi – 110034 (INDIA). Ph: +91-11-42208000, 42208111, 42208222, Mobile: +91-9810521400 Fax: +91-11-42208444 Email: customerservice@lifetechindia.com Website: www.lifetechindia.com