

Product Specification Sheet

Mouse Ceruloplasmin (Cp) Antibodies and positive control

Cat. # CP16-A	Rabbit Anti-Mouse Cp IgG, aff pure	SIZE: 100 ul
Cat. # CP16-C	Purified Mouse Cp protein control for Western Blots	SIZE: 100 ul

Ceruloplasmin (Cp) is a plasma metalloprotein synthesized and secreted by the liver. Cp (also known as ferroxidase; iron (II):oxygen oxidoreductase, EC 1.16.3.1) is a blue alpha-2-glycoprotein that binds 90 to 95% of plasma copper and has 6 or 7 cupric ions per molecule. It is involved in peroxidation of Fe(II) transferrin to form Fe(III) transferrin. Human Cp is a single polypeptide chain (1065 aa, 135 kDa; chromosome 8q21.13-q23.1) structurally similar to factors V and VIII. Cp also has abundant expression in specific populations of glial cells within the brain. Hereditary Cp deficiency have profound iron accumulation in most tissues. Cp \pm mice showed no abnormalities in cellular iron uptake but a striking impairment in the movement of iron out of reticuloendothelial cells and hepatocytes. Cp may play an essential physiologic role in determining the rate of iron efflux from cells with mobilizable iron stores.

Source of Antigen and Antibodies

Antigen	Purified Mouse Ceruloplasmin protein
Ab Host/type	Rabbit, Polyclonal purified IgG, purified over antigen-agarose (Cat # CP16-A); concn ~100 ug/100 ul
2-Ab	Goat Anti-rabbit IgG-HRP cat # 20320 (AP, biotin, FITC conjugates also available)
-ve control IgG	# 20009-1, Rabbit (non-immune) IgG, purified, suitable for ELISA, Western, IHC as -ve control

Mouse Cp protein (~120 Kda) was purified from rat serum (>95%). For Western blot +ve control (**Cat # CP16-C**) is supplied in SDS-PAGE sample buffer (reduced). Load 10 ul/lane of **CP16-C** for good visibility with antibody Cat # **CP16-A**. Store at -20oC in suitable size aliquots. SDS may crystallize in cold conditions. It should redissolve by warming before taking it from the stock. It should be heated once prior to loading on gels. If the product has been stored for several weeks, then it may be preferable to add 5 ul of fresh 2x sample buffer per 10 ul of the **CP16-C** solution prior to heating and loading on gels. This preparation is not biologically active. It is not suitable for ELISA or other applications where native protein is required. Do not freeze, thaw, or heat repeatedly

Form & Storage of Antibodies/Peptide Control

Antibody (purified)

100ug solution lyophilized powder
Supplied in Buffer: PBS, 0.2% BSA, and 0.05% azide
Reconstitute powder in 100 ul PBS

Storage

Short-term: unopened, undiluted vials for less than a week at 4oC.

Long-term: at -20C or below in suitable aliquots after reconstitution. Do not freeze and thaw and store working, diluted solutions.

Stability: 6-12 months at -20oC or below.

Shipping: 4oC for solutions and room temp for powder.

Recommended Usage

Western Blotting (1:1K-5K for aff pure IgG using Chemiluminescence technique). Cp is ~120 kDa.

ELISA (1:100K; using 50-100 ng control antigen/well).

Histochemistry & Immunofluorescence: Not tested

Specificity & Cross-reactivity

The antibodies have been tested in mouse. Antibody cross-reactivity in various species has not been studied. Antibodies to human, mouse, rat Cp and the purified proteins are also available. Purified mouse Cp protein #CP16-C can be used as positive controls for western blots.

General References: (1). Koschinsky ML et al (1986) PNAS 83, 5086-5090; Daimon M et al (1995) BBRC 208, 1028-1035; Yang F et al (1986) PNAS 83, 3257-3261; Yakahashi N et al (1984) PNAS 81, 390-394; YANG Fm et al (1990) JBC 265, 10780-10785

*This product is for in vitro research use only.

CP16-A-C

90623A

Alpha Diagnostic Intl Inc., 6203 Woodlake Center Dr, S an Antonio, T X 7 8 24 4 , U S A;

India Contact:

Life Technologies (India) Pvt. Ltd.

306, Aggarwal City Mall, Opposite M2K Pitampura, Delhi - 110034 (INDIA). Ph: +91-11-42208000, 42208111, 42208222, Mobile: +91-9810521400 Fax: +91-11-42208444 Email: customerservice@lifetechindia.com Website: www.lifetechindia.com