


Human Subcutaneous Adipose Tissue, Mesenchymal Stem Cells

Product sheet, catalog n° CTICC.1.9

General Information

- **Organism:** Human (*Homo sapiens*)
- **Tissue:** Skin
- **Cell Type:** Skin AD-MSC from single donor
- **Location:** Abdomen, face or other part (see Certificate of Analysis)
- **Gender:** Male or Female (see Certificate of Analysis)
- **Age:** (see Certificate of Analysis)
- **Phototype:** (see Certificate of Analysis)



Cell Characteristics

- **Cell properties:** Adherent
- **Morphology:** Bipolar, spindle-shaped, fibroblast-like morphology
- **Isolation:** From subcutaneous fat
- **Cell passage:** P1 to P3 (see Certificate of Analysis)
- **Minimum number of population doublings:** 15
- **Cell viability:** Minimum 80% viability when thawed from cryopreservation
- **Cell conditioning:** Supplied as vials of 1M cells
- **Cryopreservation medium:** Frozen with 90% serum-free cryopreservation medium + 10% DMSO
- **Storage condition:** Liquid nitrogen
- **Batch specific information:** included in the Certificate of Analysis

Safety and Quality Control

- **Biosafety level:** 1
- **Contamination:** Use mandatory laboratory protections and handle with care tissues and cells derived from human samples to avoid any contamination of the operator.
- **Viral testing:** Negative for HIV, HBV, HCV
- **Sterility testing:** Negative for mycoplasma, bacteria and yeasts

Handling upon delivery and storage

- Check that the containers are intact and free of damage
- If cells are not used immediately, place the vials at -150°C or below upon delivery

Growth medium

- **Recommended medium reference:** CTIGM.ADMSC: Growth Medium for Adipose Derived - Mesenchymal Stem Cells

Thawing and culturing procedure for frozen cells

- 1 - Add 0.12 ml per cm² of medium to the culture vessel (Recommended: 9mL in collagen precoated T75 flask)
- 2 - Add 13 ml of PBS solution to a 15 ml conical tube, and warm in a water bath to 37 °C
- 3 - Thaw cryovial by swirling in a water bath at 37 °C. As soon as the content has thawed, start step 4
- 4 - Once thawed, add the cell suspension to the warmed PBS in sterile conditions
- 5 - Spin the tube at 250 g for 7 minutes to pellet the cells
- 6 - Resuspend the cells in the appropriate volume of recommended medium
- 7 - Seed the cells in the culture vessel at a concentration of 5000 to 10 000 cells per cm²
- 7 - Incubate at 37°C, 5% CO₂ atmosphere, 95% humidity
- 8 - After 24 hours of incubation, change the medium to remove any debris
- 9 - Continue to incubate and change the medium every 3-4 days

Subculturing

- 1- Start subculturing when cells reached 70%-80% confluence
- 2 - Preheat TrypLE (non-toxic for cell - trypsin substitute) and recommended medium
- 3 - Remove the medium from the flask
- 4 - Wash the cells quickly with PBS without Ca²⁺ Mg²⁺
- 5 - Add 0.06mL per cm² of TrypLE for 5-15 min at 37 °C in the incubator
- 6 - Remove the cells from the flask by pipetting several times and wash the flask with recommended medium for remaining cells
- 7 - Centrifuge the cells in recommended medium at 250 g for 7 minutes to pellet the cells
- 8 - Remove the supernatant and resuspend the pellet in recommended medium
- 9 - Seed the cells in the culture vessel at a concentration of 5 000 cells per cm²
- 10 - Incubate at 37 °C, 5% CO₂ atmosphere, 95% humidity
- 11 - After 24 hours of incubation, change half of the medium to remove any debris
- 12 - Continue to incubate and change the medium every 3-4 days

Associated products

- CTICC.1.1.1: Adult Human Dermal fibroblasts, Cryopreserved, 10⁶ cells
- CTICC.1.2.1: Adult Human Epidermal Keratinocytes, Cryopreserved, 10⁶ cells
- CTICC.1.3.1: Adult Human Melanocytes, Cryopreserved, 10⁶ cells
- SKIN BIOPSIES: Fresh, Flash Frozen, FFPE, OCT-embedded

Provisions

- Cells and tissues are intended for **research use only** and shall not be used for human trials, animal trials, or diagnostics.
- **Consent:** the original tissues have been obtained after informed consent of the patient under the provisions required by French Law.
- **Primary Human cells** are not immortalised cell lines and may not be continually subcultured.