

Product Specification Sheet

Connexin 30.3 (Cx30.3)/GapJunction Beta-5 Protein (CXB5)

Cat. CX303-S	Rabbit Anti-Rat Cx30.3 Antiserum	SIZE: 100 ul
Cat. CX303-A	Rabbit Anti-Rat Cx30.3 IgG (aff Pure)	SIZE: 100 ug
Cat. CX303-P	Rat Cx30.3 Control peptide	SIZE: 100 ug

Gap junctions are composed of transmembrane channels that link the cytoplasm of neighboring cells. They differ from other membrane channels since they exist between two cells. Gap junctions are relatively non-specific and allow passive diffusion of small molecules up to 1000 Dalton. The junctions exist in almost all vertebrate and non-vertebrate cells. It is believed that gap junction play an important for intercellular communications and affect growth and differentiation of cells. Gap junctional channel is composed of a hemichannel (connexon) in the cell membrane of one cell joined in mirror symmetry with a connexon in the opposing cell. Each connexon is an oligomer of six protein subunits that define the axial aqueous pore. Molecular cloning studies have identified a family of at least 12 highly related Connexins that are designated according to mol. wt, **Cx26-50**. Hydropathy analyses of Cx sequences predicts 4 transmembraneTM, 2 extracellular (EC), and 3 cytoplasmic (CP) domains. The EC, TM, and N-terminal CP domains are well conserved among family members, while Central and C-terminal domains are highly variable in both sequence and size. The N and C-termini are predicted to be cytoplasmic.

Source of Antigen, Antibodies

Antigen	20aa peptide of Rat CX30.3 ; Designated (CX303-P or control peptide). Epitope location~ C-terminal, Cytoplasmic domain
Ab Host/type	Rabbit, polyclonal Unpurified antiserum (cat # CX303-S) Aff pure IgG (cat #CX303-A) purified over antigen-agarose column
2-ab	Goat Anti-rabbit IgG-HRP cat # 20320 (AP, biotin, FITC conjugates also available)
-ve control IgG	# 20009-1, Rabbit (non-immune) IgG, purified, suitable for ELISA, Western, IHC as -ve control

Form & Storage of Antibodies/Peptide Control

Antiserum (unpurified)

100ul solution lyophilized powder
Supplied 0.05% azide, **Reconstitute** powder in 100 ul PBS

Affinity pure IgG

100 ug/100ul solution lyophilized powder
Supplied in **Buffer:** PBS+0.1% BSA
Reconstitute powder in PBS at 1mg/ml

Control/blocking peptide

100 ug/100 ul solution lyophilized powder
Supplied in **Buffer:** PBS pH 7.5,
Reconstitute powder in PBS at 1 mg/ml.

Storage

Short-term: unopened, undiluted liquid vials at -20OC and powder at 4oC or -20oC..

Long-term: at -20C or below in suitable aliquots after reconstitution. Do not freeze and thaw and store working, diluted solutions.

Stability: 6-12 months at -20oC or below.

Shipping: 4oC for solutions and room temp for powder

Recommended Usage

Western Blotting (1:1K-5K for neat serum and 1-10 ug/ml for affinity pure using Chemiluminescence technique). The Cx30.3 has a predicted mol. Wt of of approx 31 Kda. Cx30.3 gene has been shown to be expressed in skin (see review in 1 and refs. 2-3).

ELISA (1:100K; using 50-100 ng control peptide/well).

Histochemistry & Immunofluorescence: Not Tested. We recommend the use of affinity purified antibody at 2-20 ug/ml in formaldehyde fixed tissues.

Specificity & Cross-reactivity

The rat Cx303-P immunogenic peptide sequence is identical in mouse Cx30.3 (3). It has no significant homology with other Connexins. Antibody crossreactivity with Cx30.3 in various species is not established. Control peptide, because of its low mol. Wt (<3 kDa), is not suitable for Western. It should be used for ELISA or antibody blocking experiments (use 5-10 ug control peptide per 1 ug of aff pure IgG or 1 ul antiserum) to confirm antibody specificity (see detailed protocol at the web site).

General References: (1 Kumar, Nm and Giula, NB (1996) Cell 84, 381-388; Evans, HW (1994) Biocem. Soc. Tr. 788-792; Byer, E et al (1990) J. membrane Biol. 116, 187-194; (2) Tucker, MA and Barajas, L (1994) Exp. Cell Res. 213, 224, 230; (3) Hennemann, H (1992) J. Biol. Chem. 267, 17225-17233.

2. Citations of for ADI Antibodies (see updates at the web site)

Fonseca PC, 2004, Immunol. Lett. 94, 65-75, , IHC
Laux-Fenton W, 2003, Cornea. 22(5):457-464,, , IHC

*This product is for in vitro research use only.

Related material available from ADI

Anti- Cx26-Cx50

CX303-S-A-P 71217A

Alpha Diagnostic Intl Inc., 6203 Woodlake Center Dr, S an Antonio, T X 7 8 24 4 , U S A ;

India Contact:

Life Technologies (India) Pvt. Ltd.

306, Aggarwal City Mall, Opposite M2K Pitampura, Delhi – 110034 (INDIA). Ph: +91-11-42208000, 42208111, 42208222, Mobile: +91-9810521400 Fax: +91-11-42208444 Email: customerservice@lifetechindia.com Website: www.lifetechindia.com