

Product Specification Sheet

**Connexin 43 (Cx43)/Gap Junction Alpha-1 Protein (CXA1) Antibodies**

<b>Cat.</b> CX43B12-S	Rabbit Anti-Mouse Cx43 Antiserum # 2	<b>SIZE:</b> 100 ul
<b>Cat.</b> CX43B12-A	Rabbit Anti- Mouse Cx43 IgG # 2 (aff pure)	<b>SIZE:</b> 100 ug
<b>Cat.</b> CX43B12-P	Mouse Connexin Cx43 Control/blocking peptide	<b>SIZE:</b> 100 ug

Gap junctions are composed of transmembrane channels that link the cytoplasm of neighboring cells. They differ from other membrane channels since they exist between two cells. Gap junctions are relatively non-specific and allow passive diffusion of small molecules up to 1000 Dalton. The junctions exist in almost all vertebrate and non-vertebrates cells. It is believed that gap junction play an important for intercellular communications and affect growth and differentiation of cells. Gap junctional channel is composed of a hemichannel (connexon) in the cell membrane of one cell joined in mirror symmetry with a connexon in the opposing cell. Each connexon is an oligomer of six protein subunits that define the axial aqueous pore. Molecular cloning studies have identified a family of at least 12 highly related Connexins that are designated according to mol. wt, **Cx26-50**.

**Source of Antigen, Antibodies**

<b>Antigen</b>	23-aa peptide of Mouse CX43 (gene accession # P23242, CXA1); <b>Designated (CX43B12-P or control peptide)</b> conjugated to KLH ; <b>Epitope location</b> ~ C-terminal, Cytoplasmic domain
<b>Ab Host/type</b>	Rabbit, polyclonal Unpurified antiserum (cat # CX43B12-S) Aff pure IgG (cat #CX43B12-A) purified over antigen-agarose column
<b>2-ab</b>	<b>Goat Anti-rabbit IgG-HRP</b> cat # 20320 (AP, biotin, FITC conjugates also available)
<b>-ve control</b>	# 20009-1, Rabbit (non-immune) IgG, purified, suitable for ELISA, Western, IHC as -ve control

**Form & Storage of Antibodies/Peptide Control**

**Antiserum (unpurified)**

100ul solution lyophilized powder  
Supplied 0.05% azide, **Reconstitute** powder in 100 ul PBS

**Affinity pure IgG**

100 ug/100ul solution lyophilized powder  
Supplied in **Buffer:** PBS+0.1% BSA  
**Reconstitute powder** in PBS at 1mg/ml

**Control/blocking peptide**

100 ug/100 ul solution lyophilized powder  
Supplied in **Buffer:** PBS pH 7.5,  
**Reconstitute powder** in PBS at 1 mg/ml.

**Storage**

**Short-term:** unopened, undiluted liquid vials at -200C and powder at 4oC or -20oC..

**Long-term:** at -20C or below in suitable aliquots after reconstitution. Do not freeze and thaw and store working, diluted solutions.

**Stability:** 6-12 months at -20oC or below.

**Recommended Usage**

**Western Blotting** (1:1K-5K for neat serum and 1-10 ug/ml for affinity pure using Chemiluminescence technique). See refs in 2

**ELISA** (1:100K; using 50-100 ng control peptide/well).

**Histochemistry & Immunofluorescence:** We recommend the use of affinity purified antibody at 2-20 ug/ml in formaldehyde fixed tissues. See refs in 2

**Specificity & Cross-reactivity**

Mouse CX43B12-P immunogenic peptide sequence is specific for Cx43 and no significant homology is seen with other Connexin. The mouse Cx43 peptide sequence is 100% homologous with rat, sheep and human Cx43 and 95% with chicken Cx43. Antibody crossreactivity in various other species is not established. Control peptide, because of its low mol. Wt (<3 kDa), is not suitable for Western. It should be used for ELISA or antibody blocking experiments (use 5-10 ug control peptide per 1 ug of aff pure IgG or 1 ul antiserum) to confirm antibody specificity (see detailed protocol at the web site).

**General References:** (1) Kumar, Nm and Giula, NB (1996) Cell 84, 381-388; White , WT et al (1995) Kidney Intl. 48, 1148-1157; Evans, HW (1994) Biochem. Soc. Tr. 788-792; Byer, E et al (1990) J. membrane Biol. 116, 187-194; (2) Haefliger, JA et al (1992) JBC 267, 2057-2064; ; Reed KE et al (1993) J Clin. Invest. 91, 997-1004; Wilecke K et al (1991) 114, 1049-1057; Byer EC et al (1985) JCB 105, 2621; Nicholson BJ et al (1985) JBC 260, 6514; John SA et al (1991) BBRC 178, 1312; Fishman GI et al (1990) JCB 111, 589-598.

**2. Citations for ADI Antibodies** (see updates at the web site)

Wiszniewski L, 2007, Differentiation, 75, Issue 5: 382-392, WB, Liao Y, 2007, Arterioscler. Thromb. Vasc. Biol., 27: 1037 , IHC Isakson BE, 2006, AJP Heart Circ Physiol, 290: H1199 , IHC, Sato T, 2002, Diabetes 51: 1565-1571, WB, Aase karin, 2001, Circulation 104: 358-364., , IHC, Gutstein DE, 2001, Circ. Res. 88: 333-339, WB., IHC, Theiss C, 2006, Experimental Cell Research, 313, 38-52, WB, IHC Delay RJ, 2003, Chem Senses, 28: 807 - 815, WB, Galindo EEH, 2003, Invest. Ophthalmol. Vis. Sci., 44: 2959, IHC,

\*This product is for in vitro research use only.

**Related material available from ADI**  
**Anti- Cx26 - Cx50**  
CX43B12-S-A-P 71217A

Alpha Diagnostic Intl Inc., 6203 Woodlake Center Dr, S an Antonio, T X 7 8 24 4 , U S A ;

**India Contact:**

**Life Technologies (India) Pvt. Ltd.**

306, Aggarwal City Mall, Opposite M2K Pitampura, Delhi – 110034 (INDIA). Ph: +91-11-42208000, 42208111, 42208222, Mobile: +91-9810521400 Fax: +91-11-42208444 Email: [customerservice@lifetechindia.com](mailto:customerservice@lifetechindia.com) Website: [www.lifetechindia.com](http://www.lifetechindia.com)