

Product Specification Sheet

**Dopamine D4 Receptor (D4R)**

Cat. # D4R11-P	Mouse D4 Receptor Control Peptide	<b>SIZE:</b> 100 ug
Cat. # D4R11-S	Rabbit Anti- Mouse D4 Antiserum # 1	<b>SIZE:</b> 100 ul
Cat. # D4R11-A	Rabbit Anti- Mouse D4 Ig G # 1 (affinity pure)	<b>SIZE:</b> 100 ug

Dopamine is an endogenous catecholamine that influences many cellular activities, including behavior, hormone synthesis and release, blood pressure and intracellular ion transport. A family of at least 5 Dopamine Receptors (DR) genes, D1-D5, have been identified based upon the amino acid identity, pharmacological specificity and effector responses. DR have been classified into either the D1-like (D1, D1B, and D5) or D2-like (D2, D3, and D4). The two isoforms of D2R, D2 long (D2L) and short D2S, are encoded by splice variants of a single gene and differ only by the presence of an additional 29 AA in the intracellular domain 3 of the D2 long form. It may play a role in the coupling of the receptor to G-proteins. All members of this family have similar structure and contain 7 putative transmembrane domains. A given cell or tissue may express more than one DR. Specific radioligands do not exist that can differentiate between these DR. Therefore, specific antibodies are needed to distinguish, localize, and document changes in DR levels in cells and tissues under various normal and pathological conditions.

**Source of Antigen, Antibodies**

Mouse D4 receptor is a 387 aa, G-protein coupled, and transmembrane receptor protein.

<b>Antigen</b>	10aa peptide of Mouse D4R ; <b>Designated (D4R11-P or control peptide).</b> <b>Epitope location</b> ~ Mid-region, Extracellular
<b>Ab Host/type</b>	Unpurified antiserum (cat #D4R11-S) Aff pure IgG (cat #D4R11-A)purified over the antigen column
<b>2ab</b>	Cat # 20320, goat anti-rabbit IgG-HRP (AP, biotin, FITC conjugates also available)
<b>-ve control</b>	# 20009-1, Rabbit (non-immune) IgG, purified, suitable for ELISA, Western, IHC as -ve control

**Form & Storage of Antibodies/Peptide Control**

**Antiserum (unpurified)**

100ul solution lyophilized powder  
Supplied in Buffer: 0.05% azide  
**Reconstitute** powder in 100 ul PBS

**Affinity pure IgG**

100 ug/100ul solution lyophilized powder  
Supplied in **Buffer:** PBS+0.1% BSA  
**Reconstitute powder** in PBS at 1mg/ml

**Control/blocking peptide**

100 ug/100 ul solution lyophilized powder  
Supplied in Buffer: PBS pH 7.5,  
**Reconstitute powder in PBS at 1 mg/ml.**

**Storage**

**Short-term:** unopened, undiluted liquid vials at 20°C and powder at 4°C or -20°C..

**Long-term:** at -20°C or below in suitable aliquots after reconstitution. Do not freeze and thaw and store working, diluted solutions.

**Stability:** 6-12 months at -20°C or below.

**Shipping:** 4°C for solutions and room temp for powder

**Recommended Usage**

**Western Blotting** (1:1K-5K for antiserum and 1-10 ug/ml for affinity pure IgG using Chemiluminescence technique)..

**ELISA:** Control peptide can be used to coat ELISA plates at 1 ug/ml and detected with antibodies (1:10-50K for neat serum and 0.5-1 ug/ml for affinity pure).

**Histochemistry:** not tested. We recommend the use of affinity purified antibody at 2-10 ug/ml.

**Specificity & Cross-reactivity**

10 AA mouse D4R11-P peptide showed no significant sequence homology with other dopamine receptors (D1, D2, D3 or D5). It has sequence homology of 90% with rat (385 aa) and human (467 aa) D4R receptors (1-2). Antibody crossreactivity with D4R from various species is not known. Control peptide, because of its low mol. Wt (<3 kDa), is not suitable for Western. It should be used for ELISA or antibody blocking experiments (use 5-10 ug control peptide per 1 ug of aff pure IgG or 1 ul antiserum) to confirm antibody specificity (see detailed protocol at: the web site).

**General References:**

Van Tol HH et al (1992) Nature 358, 149-152; Van Tol HH et al (1992) Nature 350, 610-614; Livingstone, CD et al (1992) Biochem. J. 287, 277-282; Seeman, P et al (1994) Am J med genet. 54, 384-390; O'Malley, KL et al (1992) New Biol. 4, 137-146; Fishburn CS et al (1995) FEBS Lett. 361, 215-219.

**Citations of ADI's antibodies for DR** (see updated list at: the web site)

\*This product is for In vitro research use only.

**Related material available from ADI**

Antibodies D1/D2/D3/D4 and Dopamine Transporter and Serotonin Transporter

**Recycle your blot in Just 5-10 min. (use the same strip for various Dopamine receptors)** New formulation will strip antibodies in just a few minutes at room temp. (no boiling or pungent mercaptoethanol).

D4R11-S-A-P

71218S

Alpha Diagnostic Intl Inc., 6203 Woodlake Center Dr, S an Antonio, T X 7 8 24 4 , U S A;

**India Contact:**

**Life Technologies (India) Pvt. Ltd.**

306, Aggarwal City Mall, Opposite M2K Pitampura, Delhi – 110034 (INDIA). Ph: +91-11-42208000, 42208111, 42208222, Mobile: +91-9810521400 Fax: +91-11-42208444 Email: [customerservice@lifetechindia.com](mailto:customerservice@lifetechindia.com) Website: [www.lifetechindia.com](http://www.lifetechindia.com)