

Product Specification Sheet

Dopamine D4 Receptor (D4R)

Cat. # D4R11-P	Mouse D4 Receptor Control Peptide	SIZE: 100 ug
Cat. # D4R11-S	Rabbit Anti- Mouse D4 Antiserum # 1	SIZE: 100 ul
Cat. # D4R11-A	Rabbit Anti- Mouse D4 Ig G # 1 (affinity pure)	SIZE: 100 ug

Dopamine is an endogenous catecholamine that influences many cellular activities, including behavior, hormone synthesis and release, blood pressure and intracellular ion transport. A family of at least 5 Dopamine Receptors (DR) genes, D1-D5, have been identified based upon the amino acid identity, pharmacological specificity and effector responses. DR have been classified into either the D1-like (D1, D1B, and D5) or D2-like (D2, D3, and D4). The two isoforms of D2R, D2 long (D2L) and short D2S), are encoded by splice variants of a single gene and differ only by the presence of an additional 29 AA in the intracellular domain 3 of the D2 long form. It may play a role in the coupling of the receptor to G-proteins. All members of this family have similar structure and contain 7 putative transmembrane domains. A given cell or tissue may express more than one DR. Specific radioligands do not exist that can differentiate between these DR. Therefore, specific antibodies are needed to distinguish, localize, and document changes in DR levels in cells and tissues under various normal and pathological conditions.

Source of Antigen, Antibodies

Mouse D4 receptor is a 387 aa, G-protein coupled, and transmembrane receptor protein.

Antigen	10aa peptide of Mouse D4R ; Designated (D4R11-P or control peptide). Epitope location ~ Mid-region, Extracellular
Ab Host/type	Unpurified antiserum (cat #D4R11-S) Aff pure IgG (cat #D4R11-A)purified over the antigen column
2ab	Cat # 20320, goat anti-rabbit IgG-HRP (AP, biotin, FITC conjugates also available)
-ve control	# 20009-1, Rabbit (non-immune) IgG, purified, suitable for ELISA, Western, IHC as -ve control

Form & Storage of Antibodies/Peptide Control

Antiserum (unpurified)

100ul solution lyophilized powder
Supplied in Buffer: 0.05% azide

Reconstitute powder in 100 ul PBS

Affinity pure IgG

100 ug/100ul solution lyophilized powder
Supplied in **Buffer:** PBS+0.1% BSA

Reconstitute powder in PBS at 1mg/ml

Control/blocking peptide

100 ug/100 ul solution lyophilized powder
Supplied in Buffer: PBS pH 7.5,

Reconstitute powder in PBS at 1 mg/ml.

Storage

Short-term: unopened, undiluted liquid vials at 20°C and powder at 4°C or -20°C..

Long-term: at -20°C or below in suitable aliquots after reconstitution. Do not freeze and thaw and store working, diluted solutions.

Stability: 6-12 months at -20°C or below.

Shipping: 4°C for solutions and room temp for powder

Recommended Usage

Western Blotting (1:1K-5K for antiserum and 1-10 ug/ml for affinity pure IgG using Chemiluminescence technique)..

ELISA: Control peptide can be used to coat ELISA plates at 1 ug/ml and detected with antibodies (1:10-50K for neat serum and 0.5-1 ug/ml for affinity pure).

Histochemistry: not tested. We recommend the use of affinity purified antibody at 2-10 ug/ml.

Specificity & Cross-reactivity

10 AA mouse D4R11-P peptide showed no significant sequence homology with other dopamine receptors (D1, D2, D3 or D5). It has sequence homology of 90% with rat (385 aa) and human (467 aa) D4R receptors (1-2). Antibody crossreactivity with D4R from various species is not known. Control peptide, because of its low mol. Wt (<3 kDa), is not suitable for Western. It should be used for ELISA or antibody blocking experiments (use 5-10 ug control peptide per 1 ug of aff pure IgG or 1 ul antiserum) to confirm antibody specificity (see detailed protocol at: the web site).

General References:

Van Tol HH et al (1992) Nature 358, 149-152; Van Tol HH et al (1992) Nature 350, 610-614; Livingstone, CD et al (1992) Biochem. J. 287, 277-282; Seeman, P et al (1994) Am J med genet. 54, 384-390; O'Malley, KL et al (1992) New Biol. 4, 137-146; Fishburn CS et al (1995) FEBS Lett. 361, 215-219.

Citations of ADI's antibodies for DR (see updated list at: the web site)

*This product is for In vitro research use only.

Related material available from ADI

Antibodies D1/D2/D3/D4 and Dopamine Transporter and Serotonin Transporter

Recycle your blot in Just 5-10 min. (use the same strip for various Dopamine receptors) New formulation will strip antibodies in just a few minutes at room temp. (no boiling or pungent mercaptoethanol).

D4R11-S-A-P

71218S

Alpha Diagnostic Intl Inc., 6203 Woodlake Center Dr, S an Antonio, T X 7 8 24 4 , U S A;

India Contact:

Life Technologies (India) Pvt. Ltd.

306, Aggarwal City Mall, Opposite M2K Pitampura, Delhi – 110034 (INDIA). Ph: +91-11-42208000, 42208111, 42208222, Mobile: +91-9810521400 Fax: +91-11-42208444 Email: customerservice@lifetechindia.com Website: www.lifetechindia.com