

Product Specification Sheet

Aromatic L-Amino Acid Decarboxylase/DOPA Decarboxylase/DDC Antibodies

| | | |
|----------------------|--|---------------------|
| Cat # DDC11-M | Mouse monoclonal Anti-Human DDC ab #1, ascites | SIZE: 100 ul |
| Cat # DDC11-P | Human DDCcontrol/blocking peptide #1 | SIZE: 100 ug |

DDC (EC 1.1.128) belongs to the family group II Decarboxylase (DDC, GAD, HDC and TYRD). DDC catalyzes the formation of tryptamine from L-tyrosine. It also acts on 5-hydroxy-LO-tryptophan and dihydroxy-L-Phenylalanine (DOPA). Human DDC (480 aa, chromosome 7p11) is a homodimer in the native state. It requires pyridoxal phosphate as a cofactor.

Source of Antigen, Antibodies

| | |
|---------------------|--|
| Antigen | 16-aa peptide of Human DDC (1) designated cat # DDC11-P. Epitope location ~ N-terminus |
| Ab Host/type | Mouse, Monoclonal (IgG2b), unpurified ascites (cat #DDC11-M) |
| 2ab | Cat # 40320, goat anti-mouse IgG-HRP (AP, biotin, FITC conjugates also available). |
| -ve control | # 20008-1, Mouse (non-immune) IgG, purified, suitable for ELISA, Western, IHC as -ve control |

Form & Storage of Antibodies/Peptide Control

Ascites (unpurified)
100ul solution lyophilized powder
Supplied in Buffer: 0.05% azide
Reconstitute powder in 100 ul PBS

Storage
Short-term: unopened, undiluted liquid vials at 20°C and powder at 4°C or -20°C..

Long-term: at -20°C or below in suitable aliquots after reconstitution. Do not freeze and thaw and store working, diluted solutions.

Stability: 6-12 months at -20°C or below.
Shipping: 4°C for solutions and room temp for powder

Recommended Usage

Western Blotting (1:1K-5K using Chemiluminescence technique). DDC is ~50 kDa.

ELISA: Control peptide can be used to coat ELISA plates at 1 ug/ml and detected with antibodies (1:10-50K).

Histochemistry: not tested.

Specificity & Cross-reactivity

The 16 AA human DDC11-P peptide shows homology with rat/ mouse/ pig (93%), and bovine/ guinea pig (87%). Antibody reacts with DDC from human, bovine, monkey, sheep, rabbit, dog, guinea pig, and rat. Crossreactivity with other species is not known. Control peptide, because of its low mol. Wt (<3 kDa), is not suitable for Western. It should be used for ELISA or antibody blocking experiments (use 5-10 ug control peptide per 1 ug of aff pure IgG or 1 ul antiserum) to confirm antibody specificity (see detailed protocol at:the web site).

General References:

Schrere LJ et al (1992) Genomics 13, 469-471; Tanaka T et al (1989) PNAS 86, 8142; Ishii, S et al (1996) J Biochem. 120, 369-376.

Citations of ADI's antibodies for DR (see updated list at: The web site)

*This product is for In vitro research use only.

Related material available from ADI

Anti-Rabbit IgG-HRP Conjugate and ECL Reagents
NuGlo Chemiluminescent Substrate Kit

Western Blot Recycling Kit (Strips blots in 5 minutes)

DDC11-M-P 71219S