

Product Specification Sheet

Estrogen Receptor β 1/ β 2 (ER β 1/2) Antibodies

Cat. ERb11-S	Rabbit Anti-Mouse ER Beta Antisera# 1	SIZE: 100 ul
Cat. ERb11-A	Rabbit Anti- Mouse ER Beta Ig G# 1 (aff pure)	SIZE: 100 ug
Cat. ERb11-P	Mouse ERb11 Control peptide	SIZE: 100 ug

Estrogens, produced by ovaries and testis, affect growth and differentiation of many target tissues. These include the male and female reproductive tissues (mammary gland, uterus, ovary, and prostate). Estrogens have also been implicated in the physiology of the bone, cardiovascular tissues, and the brain. Estrogens bind to the intracellular proteins known as estrogen receptors (ER). Estrogen receptor is a member of the super family of nuclear receptor that show a similar structure and mode of action.

Rat/mouse **ER β 1** gene encodes a protein of 485 aa with a calculated size of approx. 54 kDa. **ER β 1** is 477 aa in human. As compared to **ER α** , **ER β 1** is highly conserved in the DNA-binding domain (>90% homology) and the C-terminal ligand binding domain (55% homology). The A/B domain, the hinge region and the F-domain are not conserved. Rat ER β 1 is primarily expressed in prostate, ovary, lung, bladder, brain, uterus, and testis. ligand binding experiments with the recombinant ER β 1 revealed a single binding component for 17 β -E2 with Kd=0.6 nM. Both ER α and ER β 1 binds to common agonists and antagonists with more or less overlapping specificity. Most recently functional variants of ER β 1, termed ER β 2, have been cloned. ER β 2 has an additional 18 aa as a results of in-frame mutation within the ligand-binding domain. ER β 2 is expressed in ovary, prostate, pituitary, brain, and muscle. Variants of both ER β 1 and ER β 2 (ER β 1 δ 3, ER β 2 δ 3) were detected that show deletion of 39 aa in the DNA-binding domain. Both ER β 1 and ER β 2 specifically bind to ER response element. ER α , β 1 and ER β 2 have been found to heterodimerize with each other.

Source of Antigen and Antibodies

Antigen	16aa peptide of Mouse ER β ; Designated (ERB11-P or control peptide). epitope location ~ C-terminus
Ab Host/type	Rabbit, polyclonal Unpurified antiserum (cat #ERB11-S) Aff pure IgG (cat #ERB11-A) purified over antigen-agarose column
2-ab	Goat Anti-rabbit IgG-HRP cat # 20320 (AP, biotin, FITC conjugates also available)
-ve control IgG	# 20009-1, Rabbit (non-immune) IgG, purified, suitable for ELISA, Western, IHC as -ve control

Form & Storage of Antibodies/Peptide Control

Antiserum (unpurified)
100ul solution lyophilized powder
Supplied 0.05% azide, **Reconstitute** powder in 100 ul PBS

Affinity pure IgG
100 ug/100ul solution lyophilized powder
Supplied in **Buffer:** PBS+0.1% BSA
Reconstitute powder in PBS at 1mg/ml

Control/blocking peptide

100 ug/100 ul solution lyophilized powder
Supplied in Buffer: PBS pH 7.5,
Reconstitute powder in PBS at 1 mg/ml.

Storage

Short-term: unopened, undiluted liquid vials at -20OC and powder at 4oC or -20oC..

Long-term: at -20C or below in suitable aliquots after reconstitution. Do not freeze and thaw and store working, diluted solutions.

Stability: 6-12 months at -20oC or below.

Shipping: 4oC for solutions and room temp for powder

Recommended Usage

Western Blotting (1:1K-5K for neat serum and 1-10 ug/ml for affinity pure using Chemiluminescence technique). The predicted size of the ER β receptor is approx. 54-kDa (1).

ELISA (1:100K; using 50-100 ng ERb11 control peptide/well).

Immunoprecipitations We recommend the use of affinity pure antibodies at 2-10 ug affinity pure antibody.

Histochemistry & Immunofluorescence: Not tested. we recommend the use of affinity purified antibody at 2-20 ug/ml in formaldehyde fixed tissue.

Specificity & Cross-reactivity

The 16-aa mouse ERB11 immunogenic peptide shows 100% homology with the rat ER β 1/ER β 1 δ 3, and 87% with human ER β 1 and its variant ER β 3. The ERB11 sequence is also found in rat ER β 2 and its variant ER β 2 δ 3 (3). It is not clear, at present, if all of these ER β isoforms can be seen and distinguished by this antibody because of very small size differences in various ER β isoforms. Actual crossreactivity of antibodies in all species is not established. Control peptide, because of its low mol. Wt (<3 kDa), is not suitable for Western. It should be used for ELISA or antibody blocking experiments (use 5-10 ug control peptide per 1 ug of aff pure IgG or 1 ul antiserum) to confirm antibody specificity (see detailed protocol at the web site).

General References: Kuiper GG et al (1996) Proc Natl Acad Sci 93, 5925-5930; Tremblay, GB et al (1997) Mol. Endocrinol. 11, 353-365, Mosselman S et al (1996) FEBS Lett. 392, 49-53, Peterson DN et al (198) Endocrinology 139, 1082; Maruyama K et al (1998) BBRC 246, 142.

Citation of ADI's antibodies for ER-Beta:

Hosokawa K 2001 Mol. Hum. Reprod. 7: 137-145
Javeshghani D 2003 Hypertension, 42, 761-767 WB

*This product is for in vitro research use only.

Related material available from ADI

Monoclonal Mouse Anti-Bovine ER (crossreacts with mouse, rat, human)
ERB11-S-A-P 71217A

Alpha Diagnostic Intl Inc., 6203 Woodlake Center Dr, S an Antonio, T X 7 8 24 4 , U S A;

India Contact:

Life Technologies (India) Pvt. Ltd.

306, Aggarwal City Mall, Opposite M2K Pitampura, Delhi – 110034 (INDIA). Ph: +91-11-42208000, 42208111, 42208222, Mobile: +91-9810521400 Fax: +91-11-42208444 Email: customerservice@lifetechindia.com Website: www.lifetechindia.com