

Product Specification Sheet

GABA-A Receptor Alpha 6 Subunit (GAA6) Antibodies

Cat. # GAA16-P	Rat GAA6 control/blocking peptide # 1	SIZE: 100 ug
Cat. # GAA16-S	Rabbit Anti-Rat GAA6 antiserum # 1	SIZE: 100 ul
Cat. # GAA16-A	Rabbit Anti-Rat GAA6 IgG # 1 (aff pure)	SIZE: 100 ug

GABA (γ -amino butyric acid) is the most abundant neurotransmitter in mammalian brain. GABA exerts its effects through ionotropic ligand-gated GABA_A, GABA_C and GABA_B receptors (**GABA_BRs**). A family of GABA-A receptors subtypes exists, which are generated by alternative splicing of alpha 1-6, beta 1-4, gamma 1-4, delta, epsilon, pie, theta, and rho1-3 to form a heteromeric (pentameric?) protein complexes. Various GABA-A subunits show distinct patterns of temporal and spatial expression that may imply its tissue specific physiological role (1). **GABA A (GAA) receptor** proteins (450-627 aa) are characterized by the presence of a cleavable signal peptide, a large extracellular N-terminus, 3 TM (transmembrane) domains, a large cytoplasmic domain followed by TM4 and C-terminal extracellular domain. The regions between TM3-4 and the large cytoplasmic loop are least conserved among various GAA subunits, which may confer subunit specific functionality. GAA genes are distributed as clusters throughout the human genome (chromosomes 4, 5, 15, and X; delta subunit on chromosome 1). GAA in the brain are the targets of many clinically important drugs. Rat GAA6 protein is 443 aa protein (mouse 443 aa, human 453-aa; chromosome 5q34).

Source of Antigen and Antibodies

Antigen	13-aa peptide from rat GAA6 (1); Designation (GAA16-P, control peptide /blocking peptide) conjugated to KLH; Epitope location ~N-terminus, Extracellular domain
Antibody host/type	Rabbit, Polyclonal unpurified antiserum (Cat # GAA16-S); Rabbit, Polyclonal IgG (Cat # GAA16-A), purified over antigen-Agarose
Secondary Ab	Cat # 20320, goat anti-rabbit IgG-HRP (AP, biotin, FITC conjugates also available).
Negative Control Ab	Non-immune rabbit IgG (Cat # 20009-1) to be used as -ve control for ELISA, WB, IHC etc.

Form & Storage of Antibodies/Peptide Control

Affinity pure IgG

100 ug/100ul solution lyophilized powder
Supplied in **Buffer:** PBS+0.1% BSA
Reconstitute powder in PBS at 1mg/ml

Control/blocking peptide

100 ug/100 ul solution lyophilized powder
Supplied in Buffer: PBS pH 7.5,
Reconstitute powder in PBS at 1 mg/ml.

Storage

Short-term: unopened, undiluted vials for less than a week at 4oC.

Long-term: at -20°C or below in suitable aliquots after reconstitution. Do not freeze and thaw and store working, diluted solutions.

Stability: 6-12 months at -20°C or below.

Shipping: 4°C for solutions and room temp for powder.

Recommended Usage

Western Blotting: 1-10 μ g/ml; using affinity pure antibody (chemiluminescence technique).

ELISA: 1:100K; using 50-100 ng control peptide/well.

Histochemistry & Immunofluorescence: Not tested; we recommend the use of affinity purified antibody at 2-10 μ g/ml.

Specificity & Cross-reactivity

The GAA16-P control peptide is 100% conserved in mouse and 91% in human GAA6. No significant homology is detected with other GABA A receptor subtypes or other receptors. Antibody cross-reactivity in various species has not been studied. The **GAA16-P** Control peptide, because of its low mol. Wt (<3 kDa), is not suitable for Western. It should be used for ELISA or antibody blocking experiments (use 5-10 ug control peptide per 1 ug of aff pure IgG or 1 ul antiserum) to confirm antibody specificity (see detailed protocol at the web site)..

General References: (1). Kato K (1990) J. Mol. Biol. 214, 619-624; Luddens S (1990) nature 346, 648-651; Hicks AA (1994) Genomics 20, 285-288; Hadingham KI (1996) Mol. Pharamcol. 49, 253-259; Mehta Ak and Ticku MK (1999) Brain Res. Rev. 29, 196-271 (review);

(2) Citations of ADI's Antibodies (see web site for updated list)

Wang J (2003) "Interaction of Calcineurin and Type-A GABA Receptor γ 2 Subunits Produces Long-Term Depression at CA1 Inhibitory Synapses", Journal of Neuroscience, 23(3):826-836, WB

*This product is for In vitro research use only.

Related material available from ADI

Antibodies GABA-A receptors (α , β , and γ -subunits), GABAR associated protein, GBR1a, 1b, and GBR2, GABA

GAA16-S-A-P

71208J

India Contact:

Life Technologies (India) Pvt. Ltd.

306, Aggarwal City Mall, Opposite M2K Pitampura, Delhi - 110034 (INDIA). Ph: +91-11-42208000, 42208111, 42208222, Mobile: +91-9810521400, Fax: +91-11-42208444
Email: customerservice@lifetechindia.com Website: www.lifetechindia.com