

Product Specification Sheet

Histamine Receptor 3 (H3R) Antibodies

Cat. # H3R32-P	Human H3R Control Peptide # 2	SIZE: 100 ug
Cat. # H3R32-S	Rabbit Anti-Human H3R antiserum # 2	SIZE: 100 ul
Cat. # H3R32-A	Rabbit Anti-Human H3R Ig G # 2 (affinity pure)	SIZE: 100 ug

Histamine, one of the most important mediators of allergy and inflammation, is a chemical messenger and aminergic neurotransmitters. It plays an important role in a multitude of physiological processes in central and peripheral tissues. Histamine is synthesized in a restricted population of neurons located in the tuberomammillary nucleus of the posterior hypothalamus implicated in many brain functions (e.g. sleep/wakefulness, hormonal secretion, cardiovascular control, thermoregulation, food intake, and memory formation). In peripheral tissues histamine is stored in mast cells, basophils, enterochromaffin cells. Histamine release leads to various well-known symptoms of allergic conditions in the skin and the airway system. Histamine effects are mediated by four pharmacologically distinct receptors, the **H1R, H2R, H3R and H4R receptors**, which are all members of the G-protein coupled receptor (GPCR) family. Histamine receptors display 7 TM domains, an extracellular N-terminus, and a cytoplasmic C-terminus of variable length.

Human Histamine 3 receptor, **H3R** (GPCR97), has recently been identified as a presynaptic autoreceptor on histamine neurons in the brain controlling the stimulated release of histamine. Subsequently, the H3 receptor has been shown to be a presynaptic heteroreceptor in nonhistamine-containing neurons in both the central and peripheral nervous systems. H3R (rat/human 445 aa, ~93% homology) is primarily expressed in the brain. Several studies using H3 selective agonists revealed that H3R couples to pertussis toxin-sensitive Gi/o protein.

Source of Antigen and Antibodies

Antigen	An 18 aa peptide (Gene Accession #Q9Y5N1) (designated H3R32-P control peptide) within the extracellular, N-terminus of Human H3R (1) was synthesized, conjugated to KLH
Ab Host/type	Rabbit, Polyclonal antiserum # H3R32-S and IgG, purified over antigen-agarose (Cat # H3R32-A)
2-Ab	Cat # 20320, goat anti-rabbit IgG-HRP (AP, biotin, FITC conjugates also available).
-ve control IgG	Cat # 20009-1, Rabbit (non-immune) Serum IgG, purified, suitable for ELISA, Western, IHC as -ve control

Form & Storage of Antibodies/Peptide Control

Antiserum (unpurified)
100ul solution lyophilized powder
Supplied 0.05% azide, **Reconstitute** powder in 100 ul PBS

Affinity pure IgG
100 ug/100ul solution lyophilized powder
Supplied in **Buffer:** PBS+0.1% BSA
Reconstitute powder in PBS at 1mg/ml

Control/blocking peptide
100 ug/100 ul solution lyophilized powder
Supplied in Buffer: PBS pH 7.5,
Reconstitute powder in PBS at 1 mg/ml.

Storage
Short-term: unopened, undiluted liquid vials at -20OC and powder at 4oC or -20oC..

Long-term: at -20C or below in suitable aliquots after reconstitution. Do not freeze and thaw and store working, diluted solutions.

Stability: 6-12 months at -20oC or below.
Shipping: 4oC for solutions and room temp for powder

Recommended Usage

Western Blotting (1:1K-5K for neat serum and 1-10 ug/ml for affinity pure using Chemiluminescence technique).

ELISA (1:10K-1:100K; using 50-100 ng of control peptide/well).

Histochemistry: Not tested. We recommend the use of 2-10 ug/ml of affinity pure antibody.

Specificity & Cross-reactivity

The Human H3R32-P peptide sequence is 94% homologous to mouse, 88% conserved in rat and 84% in g. pig H3R. No significant homology exists with other histamine receptors. Antibody crossreactivity in various species is not established. Control peptide, because of its low mol. Wt (<3 kDa), is not suitable for Western. It should be used for ELISA or antibody blocking experiments (use 5-10 ug control peptide per 1 ug of aff pure IgG or 1 ul antiserum) to confirm antibody specificity see detailed protocol at the web site).

General References: Lovenberg TW et al (2000) J. Pharmacol. Exp. Ther. 293, 771-778; Lovernberg TW et al (1999) Mol. Pharmacol. 55, 1101-1107

Citations of for ADI Antibodies (see updated list at the web site)

Karlstedt k, 2003, Mol Cell Neurosci 24, 614-622, WB, IHC/cold methanol
Jokuti A, 2007, Cell Biol Int. 2007 Nov;31(11):1367-70., WB,
Sander LE, 2005, Gut 55, 498-504, WB, IF
Lippert U, 2004, J. Invest Dermatol 123, 116-123, WB,

*This product is for In vitro research use only.

Related material available from ADI
Anti-Histamine, H1R-H4R, IgE ELISA kit

H3R32-S-A-P 71214A

India Contact:

Life Technologies (India) Pvt. Ltd.

306, Aggarwal City Mall, Opposite M2K Pitampura, Delhi - 110034 (INDIA). Ph: +91-11-42208000, 42208111, 42208222, Mobile: +91-9810521400, Fax: +91-11-42208444
Email: customerservice@lifetechindia.com Website: www.lifetechindia.com