


## ML-210

|                           |   |       |          |
|---------------------------|---|-------|----------|
| <b>Cat. No.:</b>          | HY-100003   |       |          |
| <b>CAS No.:</b>           | 1360705-96-9  |       |          |
| <b>Molecular Formula:</b> | C <sub>22</sub> H <sub>20</sub> Cl <sub>2</sub> N <sub>4</sub> O <sub>4</sub> |       |          |
| <b>Molecular Weight:</b>  | 475.32  |       |          |
| <b>Target:</b>            | Glutathione Peroxidase; Ferroptosis   |       |          |
| <b>Pathway:</b>           | Metabolic Enzyme/Protease; Apoptosis  |       |          |
| <b>Storage:</b>           | Powder  | -20°C | 3 years  |
|                           |   | 4°C   | 2 years  |
|                           | In solvent  | -80°C | 6 months |
|                           |   | -20°C | 1 month  |



### SOLVENT & SOLUBILITY

|   |   |                          |              |            |            |
|---|---|--------------------------|--------------|------------|------------|
| <b>In Vitro</b>   | DMSO : 25 mg/mL (52.60 mM; Need ultrasonic)   |                          |              |            |            |
|   |   | Solvent<br>Concentration | Mass<br>1 mg | 5 mg       | 10 mg      |
|   | <b>Preparing Stock Solutions</b>  | 1 mM                     | 2.1038 mL    | 10.5192 mL | 21.0385 mL |
|   |   | 5 mM                     | 0.4208 mL    | 2.1038 mL  | 4.2077 mL  |
| 10 mM   |   | 0.2104 mL                | 1.0519 mL    | 2.1038 mL  |            |
| Please refer to the solubility information to select the appropriate solvent. |   |                          |              |            |            |
| <b>In Vivo</b>  | <ol style="list-style-type: none"> <li>Add each solvent one by one: 10% DMSO &gt;&gt; 40% PEG300 &gt;&gt; 5% Tween-80 &gt;&gt; 45% saline<br/>Solubility: 2.5 mg/mL (5.26 mM); Suspended solution; Need ultrasonic</li> <li>Add each solvent one by one: 10% DMSO &gt;&gt; 90% corn oil<br/>Solubility: ≥ 2.08 mg/mL (4.38 mM); Clear solution</li> </ol> |                          |              |            |            |

### BIOLOGICAL ACTIVITY

|                                     |  |
|-------------------------------------|--|
| <b>Description</b>                  | ML-210 is a selective and covalent glutathione peroxidase 4 (GPX4) inhibitor with an EC <sub>50</sub> of 30 nM. ML-210 binds the GPX4 selenocysteine residue. ML-210 has anti-cancer activity <sup>[1][2]</sup> .  |
| <b>IC<sub>50</sub> &amp; Target</b> | Glutathione Peroxidase 4 (GPX4) <sup>[1]</sup>   |
| <b>In Vitro</b>                     | <p>ML-210 exhibits cell-killing activity across a panel of 821 cancer cell lines (WM88, LOX-IMVI, CJM, U257, CAK12, A498, HT1080, MC38, PANC02). ML-210 is a prodrug that requires cellular activation to bind GPX4<sup>[1]</sup>.</p> <p>ML-210 has IC<sub>50</sub>s of 71 nM, 272 nM and 107nM for BJeLR (HRAS<sub>V12</sub>), BJeH-LT (without HRAS<sub>V12</sub>) and DRD cell lines, respectively<sup>[2]</sup>.</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p> |

---

## REFERENCES

- [1]. John K. Eaton, et al. Targeting a Therapy-Resistant Cancer Cell State Using Masked Electrophiles as GPX4 Inhibitors. Biorxiv. 2018.
- [2]. Weïwer M, et al. Development of small-molecule probes that selectively kill cells induced to express mutant RAS. Bioorg Med Chem Lett. 2012 Feb 15;22(4):1822-6.
- 

**Caution: Product has not been fully validated for medical applications. For research use only.**