


## AZD0156

<b>Cat. No.:</b>	HY-100016		
<b>CAS No.:</b>	1821428-35-6		
<b>Molecular Formula:</b>	C <sub>26</sub> H <sub>31</sub> N <sub>5</sub> O <sub>3</sub>		
<b>Molecular Weight:</b>	461.56		
<b>Target:</b>	ATM/ATR; Apoptosis		
<b>Pathway:</b>	Cell Cycle/DNA Damage; PI3K/Akt/mTOR; Apoptosis		
<b>Storage:</b>	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month



### SOLVENT & SOLUBILITY

<b>In Vitro</b>	DMSO : 20 mg/mL (43.33 mM; ultrasonic and adjust pH to 3 with HCl)			
		Solvent Concentration	Mass	
			1 mg	5 mg
			10 mg	
<b>Preparing Stock Solutions</b>	<b>1 mM</b>	2.1666 mL	10.8328 mL	21.6657 mL
	<b>5 mM</b>	0.4333 mL	2.1666 mL	4.3331 mL
	<b>10 mM</b>	0.2167 mL	1.0833 mL	2.1666 mL
Please refer to the solubility information to select the appropriate solvent.				
<b>In Vivo</b>	<ol style="list-style-type: none"> <li>Add each solvent one by one: 10% DMSO &gt;&gt; 40% PEG300 &gt;&gt; 5% Tween-80 &gt;&gt; 45% saline Solubility: 0.83 mg/mL (1.80 mM); Clear solution; Need ultrasonic</li> <li>Add each solvent one by one: 10% DMSO &gt;&gt; 90% (20% SBE-β-CD in saline) Solubility: 0.83 mg/mL (1.80 mM); Suspended solution; Need ultrasonic</li> <li>Add each solvent one by one: 10% DMSO &gt;&gt; 90% corn oil Solubility: 0.83 mg/mL (1.80 mM); Clear solution; Need ultrasonic</li> </ol>			

### BIOLOGICAL ACTIVITY

<b>Description</b>	AZD0156 is a potent, selective and orally active ATM inhibitor with an IC <sub>50</sub> of 0.58 nM. AZD0156 inhibits the ATM-mediated signaling, prevents DNA damage checkpoint activation, disrupts DNA damage repair, and induces tumor cell apoptosis <sup>[1]</sup> .
<b>IC<sub>50</sub> &amp; Target</b>	ATM
<b>In Vitro</b>	AZD0156 inhibits the kinase activity of ATM and ATM-mediated signaling, prevents DNA damage checkpoint activation, and disrupts DNA damage repair, induces tumor cell apoptosis, and leads to cell death in ATM-overexpressing tumor cells <sup>[1]</sup> .

---

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

---

## PROTOCOL

### Cell Assay <sup>[1]</sup>

HT29 cells are seeded into 384 well assay plates at a density of 6000 cells/well in 40 µL EMEM medium containing 1% L glutamine and 10% FBS and allowed to adhere overnight. The following morning compound of Formula (I) in 100% DMSO is added to assay plates by acoustic dispensing. After 1h incubation at 37°C and 5% CO<sub>2</sub>, 40 nL of 3 mM 4NQO in 100% DMSO is added to all wells by acoustic dispensing, except minimum control wells which are left untreated with 4NQO to generate a null response control. Plates are returned to the incubator for a further 1h. Then cells are fixed by adding 20 µL of 3.7% formaldehyde in PBS solution and incubating for 20 mins at r.t.. Then 20 µL of 0.1% Triton XI 00 in PBS is added and incubated for 10 minutes at r.t., to permeabilise cells. Then the plates are washed once with 50 µL/well PBS, using a Biotek EL405 plate washer.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

---

## CUSTOMER VALIDATION

- Cancer Discov. 2020 Nov;10(11):1742-1757.
- Cancers (Basel). 2021 Aug 20;13(16):4200.
- J Mol Med (Berl). 2019 Aug;97(8):1183-1193.

See more customer validations on [www.MedChemExpress.com](http://www.MedChemExpress.com)

## REFERENCES

[1]. Imidazo[4,5-c]quinolin-2-one compounds and their use in treating cancer.?

---

**Caution: Product has not been fully validated for medical applications. For research use only.**

India Contact:

Life Technologies (India) Pvt. Ltd.

306, Aggarwal City Mall, Opposite M2K Pitampura, Delhi – 110034 (INDIA). Ph: +91-11-42208000, 42208111, 42208222, Mobile: +91-9810521400, Fax: +91-11-42208444

Email: [customerservice@lifetechindia.com](mailto:customerservice@lifetechindia.com) Website: [www.lifetechindia.com](http://www.lifetechindia.com)