


PT2977

| | |
|---------------------------|--|
| Cat. No.: | HY-125840 |
| CAS No.: | 1672668-24-4 |
| Molecular Formula: | C ₁₇ H ₁₂ F ₃ NO ₄ S |
| Molecular Weight: | 383.34 |
| Target: | HIF/HIF Prolyl-Hydroxylase |
| Pathway: | Metabolic Enzyme/Protease |
| Storage: | -20°C, stored under nitrogen * In solvent : -80°C, 6 months; -20°C, 1 month (stored under nitrogen) |



SOLVENT & SOLUBILITY

| In Vitro | DMSO : 50 mg/mL (130.43 mM; Need ultrasonic) | | | | | | | | | | | | | | | | | | | | | | | |
|----------------------------------|---|--------------------------|------------|--|--|------|------|-------|----------------------------------|--|--|--|------|-----------|------------|------------|------|-----------|-----------|-----------|-------|-----------|-----------|-----------|
| | <table border="1"> <thead> <tr> <th rowspan="2">Solvent Concentration</th> <th colspan="3">Mass</th> </tr> <tr> <th>1 mg</th> <th>5 mg</th> <th>10 mg</th> </tr> </thead> <tbody> <tr> <td>Preparing Stock Solutions</td> <td></td> <td></td> <td></td> </tr> <tr> <td>1 mM</td> <td>2.6087 mL</td> <td>13.0433 mL</td> <td>26.0865 mL</td> </tr> <tr> <td>5 mM</td> <td>0.5217 mL</td> <td>2.6087 mL</td> <td>5.2173 mL</td> </tr> <tr> <td>10 mM</td> <td>0.2609 mL</td> <td>1.3043 mL</td> <td>2.6087 mL</td> </tr> </tbody> </table> | Solvent Concentration | Mass | | | 1 mg | 5 mg | 10 mg | Preparing Stock Solutions | | | | 1 mM | 2.6087 mL | 13.0433 mL | 26.0865 mL | 5 mM | 0.5217 mL | 2.6087 mL | 5.2173 mL | 10 mM | 0.2609 mL | 1.3043 mL | 2.6087 mL |
| Solvent Concentration | Mass | | | | | | | | | | | | | | | | | | | | | | | |
| | 1 mg | 5 mg | 10 mg | | | | | | | | | | | | | | | | | | | | | |
| Preparing Stock Solutions | | | | | | | | | | | | | | | | | | | | | | | | |
| 1 mM | 2.6087 mL | 13.0433 mL | 26.0865 mL | | | | | | | | | | | | | | | | | | | | | |
| 5 mM | 0.5217 mL | 2.6087 mL | 5.2173 mL | | | | | | | | | | | | | | | | | | | | | |
| 10 mM | 0.2609 mL | 1.3043 mL | 2.6087 mL | | | | | | | | | | | | | | | | | | | | | |
| | Please refer to the solubility information to select the appropriate solvent. | | | | | | | | | | | | | | | | | | | | | | | |
| In Vivo | <ol style="list-style-type: none"> Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.5 mg/mL (6.52 mM); Clear solution Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.5 mg/mL (6.52 mM); Clear solution Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.5 mg/mL (6.52 mM); Clear solution | | | | | | | | | | | | | | | | | | | | | | | |

BIOLOGICAL ACTIVITY

| | |
|-------------------------------------|---|
| Description | PT2977 (MK-6482) is an orally active and selective HIF-2α inhibitor with an IC ₅₀ of 9 nM. PT2977, as a second-generation HIF-2α inhibitor, increases potency and improves pharmacokinetic profile. PT2977 is a potential treatment for clear cell renal cell carcinoma (ccRCC) ^[1] . |
| IC₅₀ & Target | IC ₅₀ : 9 nM (HIF-2α) ^[1] |
| In Vitro | PT2977 potently and dose-dependently reduces mRNA levels of human cyclin D1, a target gene regulated by HIF-2α, and leads to rapid and dose-dependent reduction in EPO expression ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only. |

REFERENCES

[1]. Xu R, et al. 3-[(1S,2S,3R)-2,3-Difluoro-1-hydroxy-7-methylsulfonylindan-4-yl]oxy-5-fluorobenzonitrile (PT2977), a Hypoxia-Inducible Factor 2 α (HIF-2 α) Inhibitor for the Treatment of Clear Cell Renal Cell Carcinoma. J Med Chem. 2019 Aug 8;62(15):6876-6893.

Caution: Product has not been fully validated for medical applications. For research use only.

India Contact:

Life Technologies (India) Pvt. Ltd.

306, Aggarwal City Mall, Opposite M2K Pitampura, Delhi – 110034 (INDIA). Ph: +91-11-42208000, 42208111, 42208222, Mobile: +91-9810521400, Fax: +91-11-42208444
Email: customerservice@lifetechindia.com Website: www.lifetechindia.com