

Fucoidan

Cat. No.:	HY-132179
CAS No.:	9072-19-9
Target:	Glucosidase
Pathway:	Metabolic Enzyme/Protease
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.

Fucoidan

SOLVENT & SOLUBILITY

In Vitro	H ₂ O : 62.5 mg/mL (Need ultrasonic)
-----------------	---

BIOLOGICAL ACTIVITY

Description Fucoidan, a biologically active polysaccharide, is an efficient inhibitor of α -amylase and α -glucosidase. Anticoagulant, antitumor, antioxidant and antisteatotic activities^[1].

In Vitro Fucoidan is a biologically active polysaccharide found in great abundance in brown marine algae^[1]. Fucoidan extracted from *A. nodosum* is a more potent inhibitor of α -glucosidase, with an IC₅₀ ranging from 0.013 to 0.047 mg/mL, than the inhibition by Fucoidan extracted from *F. vesiculosus* (IC₅₀=0.049 mg/mL). Fucoidan extracted from *F. vesiculosus* does not inhibit α -amylase activity, while Fucoidan from *A. nodosum* decreases α -amylase activity by 7-100% at 5 mg/mL depending upon the algae harvest period. Fucoidan from *A. nodosum* inhibits α -amylase with an IC₅₀ of 0.12-4.64 mg/mL^[1]. Fucoidan is a fucose-rich sulfated polysaccharide typically found in the cell wall of marine algae but also recently isolated from terrestrial sources. Fucoidan exhibits an attractive therapeutic potential against a wide array of metabolic diseases associated with oxidative stress^[2]. MCE has not independently confirmed the accuracy of these methods. They are for reference only.

In Vivo Fucoidan (single injection of 25 mg/kg/body) does not inhibit tumor cell proliferation, while three-time injections of 10 mg/kg/body weight significantly reduces tumor growth and metastasis^[3]. MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Animal Model:	Mice ^[3]
Dosage:	10-400 mg/kg
Administration:	Intraperitoneal injection; followed by total-body irradiation
Result:	The mice injected with 100 mg/kg showed the best survival rate at 30 days post-irradiation.

REFERENCES

- [1]. Kyung-Tae Kim, et al. Alpha-amylase and alpha-glucosidase inhibition is differentially modulated by fucoidan obtained from *Fucus vesiculosus* and *Ascophyllum nodosum*. *Phytochemistry*. 2014 Feb;98:27-33.
- [2]. Zeinab El Rashed, et al. Antioxidant and Antisteatotic Activities of Fucoidan Fractions from Marine and Terrestrial Sources. *Molecules*. 2021 Jul 24;26(15):4467.
- [3]. Farzaneh Atashrazm, et al. Fucoidan and cancer: a multifunctional molecule with anti-tumor potential. *Mar Drugs*. 2015 Apr 14;13(4):2327-46.
-

Caution: Product has not been fully validated for medical applications. For research use only.

India Contact:

Life Technologies (India) Pvt. Ltd.

306, Aggarwal City Mall, Opposite M2K Pitampura, Delhi – 110034 (INDIA). Ph: +91-11-42208000, 42208111, 42208222, Mobile: +91-9810521400, Fax: +91-11-42208444

Email: customerservice@lifetechindia.com Website: www.lifetechindia.com