


Bedaquiline

Cat. No.:	HY-14881		
CAS No.:	843663-66-1		
Molecular Formula:	C ₃₂ H ₃₁ BrN ₂ O ₂		
Molecular Weight:	555.5		
Target:	Bacterial; Antibiotic		
Pathway:	Anti-infection		
Storage:	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month



SOLVENT & SOLUBILITY

In Vitro	DMSO : 20 mg/mL (36.00 mM; Need ultrasonic)					
		Solvent Concentration	Mass	1 mg	5 mg	10 mg
	Preparing Stock Solutions	1 mM	1.8002 mL	9.0009 mL	18.0018 mL	
		5 mM	0.3600 mL	1.8002 mL	3.6004 mL	
10 mM		0.1800 mL	0.9001 mL	1.8002 mL		
Please refer to the solubility information to select the appropriate solvent.						
In Vivo	<ol style="list-style-type: none"> Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: 2 mg/mL (3.60 mM); Suspended solution; Need ultrasonic Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: 2 mg/mL (3.60 mM); Suspended solution; Need ultrasonic Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2 mg/mL (3.60 mM); Clear solution 					

BIOLOGICAL ACTIVITY

Description	Bedaquiline (TMC207) is a diarylquinoline drug and inhibits Mycobacterium tuberculosis (Mtb) F1FO-ATP synthase through targeting of both the c- and the ε-subunit ^[1] . Bedaquiline has uncoupler activity. Bedaquiline is used for the multi-drug resistant tuberculosis ^[2] .
IC₅₀ & Target	Mtb F1FO-ATP synthase ^[1]
In Vitro	Bedaquiline inhibits the growth of TDR M. tuberculosis strains, with MIC values ranging from 0.125 to 0.5 mg/L ^[2] . Among

slowly growing mycobacteria (SGM), bedaquiline exhibits the highest activity against *Mycobacterium avium* with MIC₅₀ and MIC₉₀ values of 0.03 and 16 mg/L, respectively. Among rapidly growing mycobacteria (RGM), *Mycobacterium abscessus* subsp. *abscessus* (*M. abscessus*) and *Mycobacterium abscessus* subsp. *massiliense* (*M. massiliense*) seem more susceptible to bedaquiline than *Mycobacterium fortuitum*, with MIC₅₀ and MIC₉₀ values of 0.13 and >16 mg/L, respectively, for both species. Bedaquiline also shows moderate in vitro activity against NTM species^[3]. Bedaquiline has an excellent in vitro activity against *Mycobacterium tuberculosis*, including multidrug resistant *M. tuberculosis*^[4].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- Mbio. 2021 Jun 1;e0108821.
- Microbiol Spectr. 2021 Jun 16;e0004521.
- Antimicrob Agents Chemother. 2021 Jun 21;AAC0095621.
- Antimicrob Agents Chemother. 2021 Jan 25;AAC.01445-20.
- Antimicrob Agents Chemother. 2020 Mar 24;64(4):e02404-19.

See more customer validations on www.MedChemExpress.com

REFERENCES

- [1]. Jang JC, et al. Bedaquiline susceptibility test for totally drug-resistant tuberculosis *Mycobacterium tuberculosis*. *J Microbiol*. 2017 Apr 20.
- [2]. Pang Y, et al. In Vitro Activity of Bedaquiline against Nontuberculous Mycobacteria in China. *Antimicrob Agents Chemother*. 2017 Apr 24;61(5).
- [3]. Chahine EB, et al. Bedaquiline: a novel diarylquinoline for multidrug-resistant tuberculosis. *Ann Pharmacother*. 2014 Jan;48(1):107-15.
- [4]. Sarathy JP, et al. TBAJ-876 displays Bedaquiline-like mycobactericidal potency without retaining the parental drug's uncoupler activity. *Antimicrob Agents Chemother*. 2019 Nov 11.

Caution: Product has not been fully validated for medical applications. For research use only.

India Contact:

Life Technologies (India) Pvt. Ltd.

306, Aggarwal City Mall, Opposite M2K Pitampura, Delhi – 110034 (INDIA). Ph: +91-11-42208000, 42208111, 42208222, Mobile: +91-9810521400, Fax: +91-11-42208444

Email: customerservice@lifetechindia.com Website: www.lifetechindia.com