


## TG003

<b>Cat. No.:</b>	HY-15338		
<b>CAS No.:</b>	719277-26-6		
<b>Molecular Formula:</b>	C <sub>13</sub> H <sub>15</sub> NO <sub>2</sub> S		
<b>Molecular Weight:</b>	249.33		
<b>Target:</b>	CDK		
<b>Pathway:</b>	Cell Cycle/DNA Damage		
<b>Storage:</b>	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month



## SOLVENT & SOLUBILITY

### In Vitro

DMSO : ≥ 31 mg/mL (124.33 mM)  
 \* "≥" means soluble, but saturation unknown.

Preparing Stock Solutions	Solvent		1 mg	5 mg	10 mg
	Concentration	Mass			
	1 mM		4.0107 mL	20.0537 mL	40.1075 mL
	5 mM		0.8021 mL	4.0107 mL	8.0215 mL
	10 mM		0.4011 mL	2.0054 mL	4.0107 mL

Please refer to the solubility information to select the appropriate solvent.

### In Vivo

- Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline  
 Solubility: ≥ 2.5 mg/mL (10.03 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)  
 Solubility: ≥ 2.5 mg/mL (10.03 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% corn oil  
 Solubility: ≥ 2.5 mg/mL (10.03 mM); Clear solution

## BIOLOGICAL ACTIVITY

### Description

TG003 is a potent inhibitor of Clk1/Sty; inhibits Clk1 and Clk4 with IC<sub>50</sub> values of 20 and 15 nM, respectively<sup>[1]</sup>.

### IC<sub>50</sub> & Target

CLK1	CLK2	Clk4
20 nM (IC <sub>50</sub> )	200 nM (IC <sub>50</sub> )	15 nM (IC <sub>50</sub> )

### In Vitro

TG003, shows the most potent effect on Clk1/Sty and Clk4 (IC<sub>50</sub>, 15–20 nM) and lesser on Clk2 (200 nM). TG003 inhibits

SF2/ASF-dependent splicing of  $\beta$ -globin pre-mRNA in vitro by suppression of Clk-mediated phosphorylation. It suppresses serine/arginine-rich protein phosphorylation, dissociation of nuclear speckles, and Clk1/Sty-dependent alternative splicing in mammalian cells<sup>[1]</sup>. The small drug TG003 increases endogenous expression of p53 $\beta$  and p53 $\gamma$  protein isoforms by modulation of TP53 intron 9 alternative splicing<sup>[2]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

#### In Vivo

Intrathecal injection of either TG003 (1-100 pM) or IC261 (0.1-1 nM) dose-dependently decreases mechanical allodynia and thermal hyperalgesia induced by carrageenan or CFA<sup>[3]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

## PROTOCOL

#### Kinase Assay <sup>[1]</sup>

Kinase activity of Clks and SRPKs is assayed in a reaction mixture, containing 200 mM Tris-HCl (pH 7.5), 12.5 mM MgCl<sub>2</sub>, 8 mM dithiothreitol, 4 mM EGTA, 1–20  $\mu$ M ATP, 1  $\mu$ Ci of [ $\gamma$ -<sup>32</sup>P]ATP, 1  $\mu$ g of synthetic peptide of SF2/ASF RS domain and 0.1-1  $\mu$ g of purified kinases in a final volume of 40  $\mu$ L. The final concentration of Me<sub>2</sub>SO is adjusted to 1% regardless of inhibitor concentration. The reaction mixture is incubated at 30 or 25 °C for mammalian or Xenopus recombinant proteins, respectively, for 10 min, and a half-portion is spotted on P81 phosphocellulose membrane. The kinase assay conditions, including the incubation period and concentration of kinases and substrates, are optimized to maintain the linearity during incubation. The membrane is washed with 5% phosphoric acid solution or 5% trichloroacetic solution at least over 15 min. The radioactivity is measured using a liquid scintillation counter<sup>[1]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

#### Cell Assay <sup>[1]</sup>

2 $\times$ 10<sup>5</sup> HeLa cells or 1.5 $\times$ 10<sup>5</sup> COS-7 cells re-suspended in 2 mL of medium are plated on 6-well dishes, and 2  $\mu$ L of 10 mM TG003 dissolved in Me<sub>2</sub>SO (final concentration at 10 mM), or 2  $\mu$ L of Me<sub>2</sub>SO, is added to some wells. Cells are trypsinized, and the density is counted every 24 h for 3 days. Cells are then fixed with 1 mL of ice-cold 70% ethanol, washed with PBS, incubated in 1 mL of PBS containing 1  $\mu$ g/mL DNase-free RNase A and 50  $\mu$ g/mL propidium iodide for 20 min at 37 °C, and proceeded to cell cycle analysis<sup>[1]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

## CUSTOMER VALIDATION

- Harvard Medical School LINCS LIBRARY

See more customer validations on [www.MedChemExpress.com](http://www.MedChemExpress.com)

## REFERENCES

[1]. Muraki M, et al. Manipulation of alternative splicing by a newly developed inhibitor of Clks. *J Biol Chem*. 2004 Jun 4;279(23):24246-54.

[2]. Marcel V, et al. Modulation of p53 $\beta$  and p53 $\gamma$  expression by regulating the alternative splicing of TP53 gene modifies cellular response. *Cell Death Differ*. 2014 Sep;21(9):1377-87.

[3]. Kurihara T, et al. Alleviation of behavioral hypersensitivity in mouse models of inflammatory pain with two structurally different casein kinase 1 (CK1) inhibitors. *Mol Pain*. 2014 Mar 10;10:17.

---

**Caution: Product has not been fully validated for medical applications. For research use only.**

India Contact:  
Life Technologies (India) Pvt. Ltd.  
306, Aggarwal City Mall, Opposite M2K Pitampura, Delhi – 110034 (INDIA). Ph: +91-11-42208000, 42208111, 42208222, Mobile: +91-9810521400, Fax: +91-11-42208444  
Email: [customerservice@lifetechindia.com](mailto:customerservice@lifetechindia.com) Website: [www.lifetechindia.com](http://www.lifetechindia.com)