


Oxacillin sodium monohydrate

Cat. No.:	HY-B0465	
CAS No.:	7240-38-2	
Molecular Formula:	C ₁₉ H ₂₀ N ₃ NaO ₆ S	
Molecular Weight:	441.43	
Target:	Bacterial; Antibiotic	
Pathway:	Anti-infection	
Storage:	Powder	-20°C 3 years 4°C 2 years
	In solvent	-80°C 6 months -20°C 1 month



SOLVENT & SOLUBILITY

In Vitro

H₂O : ≥ 100 mg/mL (226.54 mM)
DMSO : 50 mg/mL (113.27 mM; Need ultrasonic)
* "≥" means soluble, but saturation unknown.

Preparing Stock Solutions	Solvent		Mass		
	Concentration		1 mg	5 mg	10 mg
	1 mM		2.2654 mL	11.3268 mL	22.6536 mL
	5 mM		0.4531 mL	2.2654 mL	4.5307 mL
	10 mM		0.2265 mL	1.1327 mL	2.2654 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

- Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline
Solubility: ≥ 2.5 mg/mL (5.66 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% corn oil
Solubility: ≥ 2.5 mg/mL (5.66 mM); Clear solution
- Add each solvent one by one: PBS
Solubility: 110 mg/mL (249.19 mM); Clear solution; Need ultrasonic

BIOLOGICAL ACTIVITY

Description

Oxacillin sodium monohydrate is an antibiotic similar to Flucloxacillin used in resistant staphylococci infections study^[1].

In Vitro

Oxacillin exhibits MIC values of ≤1 µg/mL for four mecA gene-carrying *S. aureus* clinical isolates (SA 1306, SA 1326, SA 1552, and SA 4666^[1]).
Oxacillin (5 µg/mL, 0-90 min) induces lysis of Tol⁺ and Tol⁻ strains^[2].
MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Cell Viability Assay^[2]

Cell Line:	Cells of <i>S. aureus</i> which had grown for six to eight generations in [¹⁴ C]glycerol.
Concentration:	5 µg/mL.
Incubation Time:	0, 30, 60, 90 min.
Result:	The Tol ⁺ isolates were readily distinguished from Tol ⁻ isolates by the rates at which the cells lysed.

CUSTOMER VALIDATION

- Biomed Res Int. 2018 Jul 2;2018:3579832.

See more customer validations on www.MedChemExpress.com

REFERENCES

- [1]. Alexandros Ikonmidis, et al. In vitro and in vivo evaluations of oxacillin efficiency against mecA-positive oxacillin-susceptible *Staphylococcus aureus*. *Antimicrob Agents Chemother*. 2008 Nov;52(11):3905-8.
- [2]. R H Raynor, et al. Oxacillin-induced lysis of *Staphylococcus aureus*. *Antimicrob Agents Chemother*. 1979 Aug;16(2):134-40.

Caution: Product has not been fully validated for medical applications. For research use only.

India Contact:
Life Technologies (India) Pvt. Ltd.
306, Aggarwal City Mall, Opposite M2K Pitampura, Delhi – 110034 (INDIA). Ph: +91-11-42208000, 42208111, 42208222, Mobile: +91-9810521400, Fax: +91-11-42208444
Email: customerservice@lifetechindia.com Website: www.lifetechindia.com