


Glycyrrhizic acid

| | | | |
|---------------------------|---|-------|----------|
| Cat. No.: | HY-N0184 | | |
| CAS No.: | 1405-86-3 | | |
| Molecular Formula: | C ₄₂ H ₆₂ O ₁₆ | | |
| Molecular Weight: | 822.93 | | |
| Target: | Virus Protease | | |
| Pathway: | Anti-infection | | |
| Storage: | Powder | -20°C | 3 years |
| | | 4°C | 2 years |
| | In solvent | -80°C | 6 months |
| | | -20°C | 1 month |



SOLVENT & SOLUBILITY

In Vitro

DMSO : 500 mg/mL (607.59 mM; Need ultrasonic)
 H₂O : < 0.1 mg/mL (insoluble)

| Preparing Stock Solutions | Solvent / Mass | | 1 mg | 5 mg | 10 mg |
|---------------------------|----------------|------|-----------|-----------|------------|
| | Concentration | Mass | | | |
| Preparing Stock Solutions | 1 mM | | 1.2152 mL | 6.0759 mL | 12.1517 mL |
| | 5 mM | | 0.2430 mL | 1.2152 mL | 2.4303 mL |
| | 10 mM | | 0.1215 mL | 0.6076 mL | 1.2152 mL |

Please refer to the solubility information to select the appropriate solvent.

In Vivo

- Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline
 Solubility: ≥ 22.5 mg/mL (27.34 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)
 Solubility: 10 mg/mL (12.15 mM); Suspended solution; Need ultrasonic
- Add each solvent one by one: 10% DMSO >> 90% corn oil
 Solubility: 10 mg/mL (12.15 mM); Suspended solution; Need ultrasonic

BIOLOGICAL ACTIVITY

Description

Glycyrrhizic acid is a triterpenoid saponin, acting as a direct HMGB1 antagonist, with anti-tumor, anti-diabetic activities.

In Vitro

Glycyrrhizic acid shows a series of anti-cancer-related pharmacological activities, such as broad-spectrum anti-cancer ability, resistance to the tissue toxicity caused by chemotherapy and radiation, drug absorption enhancing effects and anti-multidrug resistance (MDR) mechanisms, as a carrier material in drug delivery systems^[1].
 In intestinal NCI-H716 cells that secretes GLP-1, Glycyrrhizic acid promotes GLP-1 secretion with a marked elevation of calcium levels. Glycyrrhizic acid can enhance GLP-1 secretion through TGR5 activation^[2].

Glycyrrhizic acid can form a stable transparent low-molecular-weight hydrogels (LMWHs) at 37°C in physiological phosphate buffered saline (PBS) with nanoclusters as the microstructures^[4].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

In Vivo

In type 1-like diabetic rats induced by streptozotocin (STZ-treated rats), Glycyrrhizic acid increases the level of plasma GLP-1, which is blocked by triamterene at a dose sufficient to inhibit Takeda G-protein-coupled receptor 5 (TGR5)^[1]. Glycyrrhizic acid (Glycyrrhizic acid; 50 mg/kg, i.p.) significantly decreases the levels of TgAb, HMGB1, TNF- α , IL-6, IL-1 β in mice^[3].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

PROTOCOL

Animal Administration ^[3]

Mice^[3]

NOD.H-2^{h4} mice are fed in the animal house until 4 weeks of age. A total of 24 male mice are then randomly separated into three different groups (n=8 per group). Mice in the control group are given sterile water without supplement. Mice in the iodine supplement (NaI) group are given 0.005% NaI in the drinking water. Mice in the NaI+Glycyrrhizic acid group are treated with 50 mg/kg Glycyrrhizic acid once daily for 4 weeks, administered via intraperitoneal injection after 8 weeks of iodine supplementation. Thyroid tissues are removed surgically under anesthesia, washed with cold saline, blotted on filter paper and weighed using an electronic balance. The thyroid gland tissues are then stored at -80°C until use^[3].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- Cell Prolif. 2020 Jun;53(6):e12829.
- Front Cell Dev Biol. 2020 Aug 11;8:713.
- Aging (Albany NY). 2019 Jul 7;11(13):4323-4337.
- ACS Chem Neurosci. 2020 Mar 18;11(6):979-993.
- J Funct Foods. 2021, 104584.

See more customer validations on www.MedChemExpress.com

REFERENCES

[1]. Su X, et al. Glycyrrhizic acid: A promising carrier material for anticancer therapy. Biomed Pharmacother. 2017 Sep 5;95:670-678

[2]. Wang LY, et al. Glycyrrhizic acid increases glucagon like peptide-1 secretion via TGR5 activation in type 1-like diabetic rats. Biomed Pharmacother. 2017 Sep 4;95:599-604.

[3]. Li C, et al. Glycyrrhizin, a Direct HMGB1 Antagonist, Ameliorates Inflammatory Infiltration in a Model of Autoimmune Thyroiditis via Inhibition of TLR2-HMGB1 Signaling. Thyroid. 2017 May;27(5):722-731.

[4]. Xia Zhao, et al. A Simple Injectable Moldable Hydrogel Assembled from Natural Glycyrrhizic Acid with Inherent Antibacterial Activity. ACS Appl. Bio Mater. 2020, 3, 1, 648-653.

Caution: Product has not been fully validated for medical applications. For research use only.

India Contact:
Life Technologies (India) Pvt. Ltd.
306, Aggarwal City Mall, Opposite M2K Pitampura, Delhi – 110034 (INDIA). Ph: +91-11-42208000, 42208111, 42208222, Mobile: +91-9810521400, Fax: +91-11-42208444
Email: customerservice@lifetechindia.com Website: www.lifetechindia.com