

Overview

Synonyms	BD 2 Murine ; BD-2 Murine
Description	Defensins (alpha and beta) are cationic peptides with a broad spectrum of antimicrobial activity that comprise an important arm of the innate immune system. The α -defensins are distinguished from the β -defensins by the pairing of their three disulfide bonds. To date, four β -defensins have been identified; BD-1, BD-2, BD-3 and BD-4. β -defensins are expressed on some leukocytes and at epithelial surfaces. In addition to their direct antimicrobial activities, they are chemoattractant towards immature dendritic cells and memory T cells. The β -defensin proteins are expressed as the C-terminal portion of precursors and are released by proteolytic cleavage of a signal sequence and, in the case of BD-1 (36 a.a.), a propeptide region. β -defensins contain a six-cysteine motif that forms three intra-molecular disulfide bonds.
Species	Mouse
Source	<i>E. coli</i>
Biological Activity	Fully biologically active when compared to standard. The biologically active determined by a chemotaxis bioassay using immature human dendritic cells is in a concentration of 10-100 ng/ml.
Sequence	AVGSLKSIQY EAELDHCHTN GGYCVRAICP PSARRPGSCF PEKNPCKYK K

Properties

Measured Molecular Weight	Approximately 5.5 kDa, a single non-glycosylated polypeptide chain containing 51 amino acids.
Purity	> 98 % by SDS-PAGE and HPLC analyses.
Formulation	Lyophilized from a 0.2 μ m filtered concentrated solution in PBS, pH 7.4.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at -20 °C. Further dilutions should be made in appropriate buffered solutions.
Endotoxin Level	Less than 1 EU/ μ g of rMuBD-2 as determined by LAL method.
Physical Appearance	Sterile Filtered White lyophilized (freeze-dried) powder.
Storage	This lyophilized preparation is stable at 2-8 °C, but should be kept at -20 °C for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one week at 2-8 °C. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20 °C to -70 °C. Avoid repeated freeze/thaw cycles.

India Contact:

Life Technologies (India) Pvt. Ltd.

306, Aggarwal City Mall, Opposite M2K Pitampura,
Delhi – 110034 (INDIA).

Mobile: +91-9810521400, Ph: +91-11-42208000

Email: customerservice@lifetechindia.com

Web: www.lifetechindia.com