

Product Specification Sheet

Human mutY Antibodies

Cat # MUTY12-P	Human mutY Control/blocking peptide #2	SIZE: 100 ug
Cat # MUTY12-S	Rabbit Anti-Human mutY antiserum # 2	SIZE: 100 ul
Cat # MUTY12-A	Rabbit Anti-Human mutY IgG # 2, aff pure	SIZE: 100 ug

Escherichia coli **MutY** is an adenine DNA glycosylase active on DNA substrates containing A/G, A/8-oxoG, or A/C mismatches and also has a weak guanine glycosylase activity on G/8-oxoG-containing DNA. The N-terminal domain of MutY, residues 1-226, has been shown to retain catalytic activity. The post-replicative adenine:8-oxo-7,8-dihydroguanine (GO) mismatch is crucial for G:C to T:A transversion. This mismatch is corrected by Escherichia coli MutY which excises the adenine from A:GO. A candidate gene coding for the human counterpart of MutY has been cloned as hMYH (535 aa) (1).

Source of Antigen and Antibodies

Antigen	19-aa peptide from human mutY-(1) ; Designation (#MUTY12-P, control/blocking peptide) conjugated to KLH; epitope location ~ C-terminus
Ab Host/type	Rabbit, Polyclonal unpurified antiserum (#MUTY12-S) and IgG, purified over antigen-agarose (Cat # MUTY12-A)
2-Ab	Cat # 20320, goat anti-rabbit IgG-HRP (AP, biotin, FITC conjugates also available).
-ve control IgG	# 20009-1, Rabbit (non-immune) IgG, purified, suitable for ELISA, Western, IHC as -ve control

Form & Storage of Antibodies/Peptide Control

Antiserum (unpurified)

100ul solution lyophilized powder
Supplied 0.05% azide, **Reconstitute** powder in 100 ul PBS

Affinity pure IgG

100 ug/100ul solution lyophilized powder
Supplied in **Buffer:** PBS+0.1% BSA
Reconstitute powder in PBS at 1mg/ml

Control/blocking peptide

100 ug/100 ul solution lyophilized powder
Supplied in **Buffer:** PBS pH 7.5,
Reconstitute powder in PBS at 1 mg/ml.

Storage

Short-term: unopened, undiluted liquid vials at -20OC and powder at 4oC or -20oC..

Long-term: at -20C or below in suitable aliquots after reconstitution. Do not freeze and thaw and store working, diluted solutions.

Stability: 6-12 months at -20oC or below.

Shipping: 4oC for solutions and room temp for powder

Recommended Usage

Western Blotting: An initial dilution of 1:1K-5K is recommended for Western. Human mutY is approx. 55 kDa protein. Users must optimize antibody dilution depending upon the nature of samples and other technical conditions.

ELISA (1:10-50K; using 50-100 ng control peptide/well).

Histochemistry & Immunofluorescence: not tested. An initial dilution of 1:500:1K is recommended for IHC. Users must optimize antibody dilution depending upon the nature of samples and other technical conditions.

Specificity & Cross-reactivity

Human MUTY12-P peptide sequence has no significant homology with E. coli mutY (1). Antibody crossreactivity in various species is not known. **Control peptide**, because of its low mol size (<3 Kda), is not recommended for Western. It should be used in ELISA, dot blot, and for antibody blocking to confirm specificity of antibodies.

General References: 1. Slupska MM et al (1996) J. Bacteriol. 178, 3885-3892; Slupska MM et al (1999) J. Bacteriol 181, 6210-3 Ohtsubo T, et al (2000) Nucleic Acids Res 15:1355-1364; Wright PM et al (1999) J Biol Chem 274, 29011-29018; Takao M et al (1999) Nucleic Acids Res. 27, 3638-44; Noll DM et al (1999) Biochemistry;38, 6374-9

***This product is for In vitro research use only.**

Related material available from ADI

Antibodies to MGMT, hNTH, Ape, mutY, OGG1, XRCC1, DNA Ligase, Polymerase, XPG, ORC1 etc are available.

MUTY12-S-A-P

71205A