

Product Specification Sheet

**NEPH1 Antibodies**

Cat. # NEPH11-P Rat NEPH1 control/blocking peptide # 1 **SIZE:** 100 ug

Cat. # NEPH11-A **Rabbit Anti-Rat NEPH1 IgG # 1 (aff pure)** **SIZE:** 100 ug

**Nephrin** is encoded by NPHS1 gene (chromosome 19q13.1), the core protein of the interpodocytes slit diaphragm of kidney glomerulus, these podocytes are highly differentiated with characteristic interdigitating foot processes covering the outer glomerular basement membrane, the space between these foot processes is spanned by a tight junction to provide the large surface area for filtration.

The mutated NPHS1 gene in congenital nephrotic syndrome of Finnish type, a human disease that leads to massive proteinuria in utero and nephrosis at birth. Nephrin is a transmembrane protein of immunoglobulin super family with 8 Ig like domains, identified in human (1241aa), rat (1252aa), and mouse (1256aa). Nephrin is found to be specifically expressed in kidney-slit diaphragm, brain and pancreas, the protein plays a crucial role in development and function of kidney filtration barrier, this barrier is crucial for maintaining the water and electrolyte balance without losing circulating proteins into the urine.

Nephrin interacts with podocin and with CD2AP C-terminal domain, It tends to be a signaling molecule that activates canonical protein kinase cascades, which is initiated by three closely related proteins called NEPH proteins (NEPH1, NEPH2, & NEPH3), The NEPH proteins share a common domain architecture consisting of 5 extracellular Ig domain followed by transmembrane domain, These proteins bind to C-terminal domain of podocin, which interacts with C-terminal of nephrin and greatly enhances nephrin-induced signaling.

**NEPH1** (Human 605aa, 67kD protein; Mouse 789aa, the sequence is 42.96% conserved between human and mouse). NEPH1 was identified in mice by a retrovirus mediated mutagenesis screening, contains 5 Ig like domains and is structurally related to nephrin. The protein has a putative signal sequence at N-terminus and a transmembrane region. Inactivation of NEPH1 develops a nephrotic syndrome that resembles NPHS mutations, suggesting that all three proteins are essential for the integrity of glomerular podocytes. NEPH1 is expressed in kidney, brain, and smooth muscle of liver, lung and heart.

**FUNCTION:** Plays a significant role in the normal development and function of the glomerular permeability

**SUBCELLULAR LOCATION:** Membrane; Single-pass type I membrane protein

**SIMILARITY:** Belongs to the immunoglobulin superfamily.

**Protein name** Kin of IRRE-like protein 1 [Precursor]

**Synonyms** Kin of irregular chiasm-like protein 1

Nephrin-like protein 1

**Gene name** Name: Kirrel1 **Synonyms:** Kirrel, Neph1

**Source of Antigen and Antibodies**

<b>Antigen</b>	16-aa peptide from Rat NEPH1;(protein accession #Q6X936, refs 1) <b>Designation (NEPH11-P, control peptide)</b> conjugated to KLH; Epitope location ~C-terminus, Cytoplasmic domain
<b>Ab Host/type</b>	Rabbit, Polyclonal Aff pure IgG (cat # NEPH11-A) purified over the antigen column
<b>2-ab</b>	Cat # 20320, goat anti-rabbit IgG-HRP (AP, biotin, FITC conjugates also available)
<b>-ve control</b>	# 20009-1, Rabbit (non-immune) IgG, purified, suitable for ELISA, Western, IHC as -ve control

**Form & Storage of Antibodies/Peptide Control**

**Affinity pure IgG**

100 ug/100ul solution lyophilized powder  
Supplied in **Buffer:** PBS+0.1% BSA  
**Reconstitute powder** in PBS at 1mg/ml

**Control/blocking peptide**

100 ug/100 ul solution lyophilized powder  
Supplied in **Buffer:** PBS pH 7.5,  
**Reconstitute powder in PBS at 1 mg/ml.**

**Storage**

**Short-term:** unopened, undiluted liquid vials at -20OC and powder at 4oC or -20oC..

**Long-term:** at -20C or below in suitable aliquots after reconstitution. Do not freeze and thaw and store working, diluted solutions.

**Stability:** 6-12 months at -20oC or below.

**Shipping:** 4oC for solutions and room temp for powder

**Recommended Usage**

**Western Blotting** (1-10 ug/ml for affinity pure antibody using ECL technique).

**ELISA:** Control peptide can be used to coat ELISA plates at 1 ug/ml and detected with antibodies (0.5-1 ug/ml for affinity pure).

**Histochemistry & Immunofluorescence:** Not tested. We recommend the use of aff pure IgG at 2-20 ug/ml.

**Specificity & Cross-reactivity**

The Rat NEPH11 control peptide is 100% conserved in human and mouse. No significant sequence homology is seen with other related family members Antibody cross-reactivity in various species has not been studied. Control peptide, because of its low mol. Wt (<3 kDa), is not suitable for Western. It should be used for ELISA or antibody blocking experiments (use 5-10 ug control peptide per 1 ug of aff pure IgG or 1 ul antiserum) to confirm antibody specificity (see detailed protocol at: [www.4adi.com/data/abblock.html](http://www.4adi.com/data/abblock.html)).

**General References:** Lorenz Sellin et al (2002) FASEB journal article 10.1096, published online; Heli Putaala et al (2001) Human Mol. Gen.10, 1-8; Dorbit B.Donoviel et al (2001) Mol and Cellular Biol 21, 4829-4836, David B. Kershaw et al (1997) JBC 272, 15708-15714.

\*This product is for In vitro research use only.

**Related material available from ADI**

Antibodies & Peptides: Nephrin related proteins (NEPH 1-3, Filtrin, Podocin, Podocalyxin).

NEPH11-A-P 70828J

**India Contact:**

**Life Technologies (India) Pvt. Ltd.**

306, Aggarwal City Mall, Opposite M2K Pitampura, Delhi - 110034 (INDIA). Ph: +91-11-42208000, 42208111, 42208222, Mobile: +91-9810521400, Fax: +91-11-42208444

Email: [customerservice@lifetechindia.com](mailto:customerservice@lifetechindia.com) Website: [www.lifetechindia.com](http://www.lifetechindia.com)