

Product Specification Sheet

Na⁺/H⁺ Exchanger 2 (NHE2) Antibodies

Cat. NHE22-S	Chicken Anti-Rat NHE2 (Antiserum # 2)	SIZE: 100 ul
Cat. NHE22-A	Chicken Anti- Rat NHE2 (affinity pure) Ig G # 2	SIZE: 100 ug
Cat. NHE22-P	Rat NHE2 control/blocking peptide	SIZE: 100 ug

Na⁺/H⁺ exchangers (**NHE**) of mammalian cells are plasma membrane intrinsic proteins mediating exchange of N⁺ and H⁺ ions in various tissues. The NHE catalyzes the electroneutral transport of extracellular Na⁺ for intracellular H⁺. They play a major role in regulation of intracellular pH (pHi) in addition to trans-cellular absorption of Na⁺, cell volume regulation and possibly in cell proliferation. These primary functions of the Na⁺/H⁺ exchanger have been related to many pathophysiological states, include hypertension, organ growth and hypertrophy, regression of cancer and renal intestinal disorders. Five NHE isoforms (NHE1-5) have been cloned so far. They are all similar in their primary structure and predicted to have 10-12 transmembrane domains. The COOH-terminals of NHE1, NHE2 and NHE3 are intracellular.

NHE2 (rat 813 aa; human chromosome 2 and rat chromosome 9) has been implicated in volume regulation in renal inner medullary collecting duct cells. Its mRNA is found in kidney medulla, cortex, colon, jejunum, ileum, human jejunum, ileum, duodenum, stomach and adrenal glands (1).

FUNCTION: Involved in pH regulation to eliminate acids generated by active metabolism or to counter adverse environmental conditions. Major proton extruding system driven by the inward sodium ion chemical gradient. Seems to play an important role in colonic sodium absorption.

SUBCELLULAR LOCATION: Membrane; Multi-pass membrane protein

SIMILARITY: Belongs to the Na(+)/H(+) exchanger (TC 2.A.36.5.1) family [view classification].

Protein name Sodium/hydrogen exchanger 2

Synonyms Na(+)/H(+) exchanger 2

NHE-2, Solute carrier family 9 member 2, H7

Gene name Name: Slc9a2 Synonyms: Nhe2

Source of Antigen and Antibodies

Antigen	20aa peptide of Rat NHE2/Slc9a2 ; (protein accession #P48763 , refs 1) Designated (NHE22-P or control peptide) , conjugated to KLH; Epitope location~C-terminal, Cytoplasmic
Ab Host/type	Chicken, polyclonal Unpurified antiserum (cat #NHE22-S) Aff pure IgG (cat #NHE22-A) purified over the antigen column
Ab Format	Cat # 60320, goat anti-chicken IgG-HRP (AP, biotin, FITC conjugates also available
-ve control	# 20010-1, Chicken (non-immune) IgG, purified, suitable for ELISA, Western, IHC as -ve control

Form & Storage of Antibodies/Peptide Control

Antiserum (unpurified)

100ul solution lyophilized powder

Supplied in Buffer: 0.05% azide

Reconstitute powder in 100 ul PBS

Affinity pure IgG

100 ug/100ul solution lyophilized powder

Supplied in **Buffer:** PBS+0.1% BSA

Reconstitute powder in PBS at 1 mg/ml

Control/blocking peptide

100 ug/100 ul solution lyophilized powder

Supplied in Buffer: PBS pH 7.5,

Reconstitute powder in PBS at 1 mg/ml.

Storage

Short-term: unopened, undiluted liquid vials at -20OC and powder at 4oC or -20oC..

Long-term: at -20C or below in suitable aliquots after reconstitution. Do not freeze and thaw and store working, diluted solutions.

Stability: 6-12 months at -20oC or below.

Recommended Usage

Western Blotting (1:1K-5K for neat serum and 1-10 ug/ml for affinity pure using Chemiluminescence technique). NHE2 is ~90 kDa (2).

ELISA (1:100K; using 50-100 ng control peptide/well).

Histochemistry & Immunofluorescence: Not tested. we recommend the use of affinity purified antibody at 2-20 ug/ml in formaldehyde fixed tissue.

Specificity & Cross-reactivity

The 20-aa rat NHE22-P sequence shows 100% homology with human, and 85% with the rabbit NHE2. No significant sequence homology exist with other NHE isoforms (NHE1, 3-5). Actual crossreactivity of antibodies in all species is not established. Control peptide, because of its low mol. Wt (<3 kDa), is not suitable for Western. It should be used for ELISA or antibody blocking experiments (use 5-10 ug control peptide per 1 ug of aff pure IgG or 1 ul antiserum) to confirm antibody specificity (see detailed protocol at: www.4adi.com/data/abblock.html).

General References:

Wang Z et al (1993) J Biol Chem. 268, 11925; Collins JF et al (1990) PNAS 90, 9398; Ghishan FK et al (1995) Genomics 30, 25; Chris Yun CH et al (1995) Am J Physiol. 269, G1-G11 (Review); Josette N and Pouyssegur J (1995) Am J Physiol. 268, C283-C296 (review); Brookstein C et al (1994) J Biol. Chem. 269, 29704.

Citations of ADI's antibodies for NHE (see updated list at: www.4adi.com/flr/nheflr.html)

*This product is for in vitro research use only.

Some New Antibodies from ADI...

• **Neurotransmitter Transporters** • Acetylcholine • BGT-1 • Dopamine • GABA (GAT1, -2, -3) • VGAT • Glycine (Glyt 1, 2) • Glutamate (GLT1, GLAST, EAAC1) • Serotonin • Proline, Vesicular GABA Transporter (VGAT) • NET • Taurine • **Renal Physiology** Arginine Vasopressin Receptors 1/2 • Aquaporins (AQP 1-5) and Urea transporter (RUT2)

NHE22-S-A-P

70829J

India Contact:

Life Technologies (India) Pvt. Ltd.

306, Aggarwal City Mall, Opposite M2K Pitampura, Delhi – 110034 (INDIA). Ph: +91-11-42208000, 42208111, 42208222, Mobile: +91-9810521400, Fax: +91-11-42208444

Email: customerservice@lifetechindia.com Website: www.lifetechindia.com