

Product Specification Sheet

Na⁺/H⁺ Exchanger 4 (NHE4) Antibodies

Cat. NHE41-S	Rabbit Anti-Rat NHE4 (Antiserum # 1)	SIZE: 100 ul
Cat. NHE41-A	Rabbit Anti- Rat NHE4 (affinity pure) Ig G # 1	SIZE: 100 ug
Cat. NHE41-P	Rat NHE4 control/blocking peptide	SIZE: 100 ug

Na⁺/H⁺ exchangers (**NHE**) of mammalian cells are plasma membrane intrinsic proteins mediating exchange of N⁺ and H⁺ ions in various tissues. The NHE catalyzes the electroneutral transport of extracellular Na⁺ for intracellular H⁺. They play a major role in regulation of intracellular pH (pHi) in addition to trans-cellular absorption of Na⁺, cell volume regulation and possibly in cell proliferation. These primary functions of the Na⁺/H⁺ exchanger have been related to many pathophysiological states, include hypertension, organ growth and hypertrophy, regression of cancer and renal intestinal disorders. Five NHE isoforms (NHE1-5) have been cloned so far. They are all similar in their primary structure and predicted to have 10-12 transmembrane domains. The COOH-terminals of NHE1, NHE2 and NHE3 are intracellular.

NHE4 (rat, 717aa; rat chromosome 9 and human chromosome 2) expression is most abundant in rat stomach followed by colon and intestine, and in low levels in rat kidney, uterus, and brain (1). Unlike other NHEs, NHE4 has no amiloride-sensitive Na⁺ uptake under basal or acid-loaded isotonic conditions.

FUNCTION: Involved in pH regulation to eliminate acids generated by active metabolism or to counter adverse environmental conditions. Major proton extruding system driven by the inward sodium ion chemical gradient. Plays an important role in signal transduction.

SUBCELLULAR LOCATION: Membrane; Multi-pass membrane protein.

SIMILARITY: Belongs to the Na(+)/H(+) exchanger (TC 2.A.36.5.1) family [view classification].

Protein name Sodium/hydrogen exchanger 4

Synonyms Na(+)/H(+) exchanger 4, NHE-4

Solute carrier family 9 member 4

Gene name Name: Slc9a4; Synonyms: Nhe4

Source of Antigen and Antibodies

Antigen	17aa peptide of Rat NHE4/Slc9a4 ; (gene accession #P26434) Designated (NHE41-P or control peptide) . conjugated to KLH, Epitope location ~C-terminal. Cytoplasmic
Ab Host/type	Rabbit, polyclonal; Unpurified antiserum (cat #NHE41-S) Aff pure IgG (cat #NHE41-A) purified over the antigen column
2-ab	Goat Anti-rabbit IgG-HRP cat # 20320 (AP, biotin, FITC conjugates also available)
-ve control	# 20009-1, Rabbit (non-immune) IgG, purified, suitable for ELISA, Western, IHC as -ve control

Form & Storage of Antibodies/Peptide Control

Antiserum (unpurified)

100ul solution lyophilized powder
Supplied in Buffer: 0.05% azide
Reconstitute powder in 100 ul PBS

Affinity pure IgG

100 ug/100ul solution lyophilized powder
Supplied in **Buffer:** PBS+0.1% BSA
Reconstitute powder in PBS at 1mg/ml

Control/blocking peptide

100 ug/100 ul solution lyophilized powder
Supplied in Buffer: PBS pH 7.5,

Reconstitute powder in PBS at 1 mg/ml.

Storage

Short-term: unopened, undiluted liquid vials at -20OC and powder at 4oC or -20oC..

Long-term: at -20C or below in suitable aliquots after reconstitution. Do not freeze and thaw and store working, diluted solutions.

Stability: 6-12 months at -20oC or below.

Recommended Usage

Western Blotting (1:1K-5K for neat serum and 1-10 ug/ml for affinity pure using Chemiluminescence technique).

ELISA (1:100K; using 50-100 ng control peptide/well).

Histochemistry & Immunofluorescence: Not tested. we recommend the use of affinity purified antibody at 2-20 ug/ml in formaldehyde fixed tissue.

Specificity & Cross-reactivity

The 17-aa rat NHE41-P sequence is unique to rat NHE4. No significant sequence homology exist with other NHE isoforms (NHE1, 2, 3, and 5). NHE4 from other species are yet to be cloned. Actual crossreactivity of antibodies in various species is not established. Control peptide, because of its low mol. Wt (<3 kDa), is not suitable for Western. It should be used for ELISA or antibody blocking experiments (use 5-10 ug control peptide per 1 ug of aff pure IgG or 1 ul antiserum) to confirm antibody specificity (see detailed protocol at: www.4adi.com/data/abblock.html).

General References:

Orlowski J et al (1992) J Biol. Chem. 267, 9331; Chris Yun CH et al (1995) Am J Physiol. 269, G1-G11 (Review); Josette N and Pouyssegur J (1995) Am J Physiol. 268, C283-C296 (review).

Citations of ADI's antibodies for NHE (see updated list at: www.4adi.com/flr/nheflr.html)

*This product is for in vitro research use only.

Some New Antibodies from ADI...

• **Neurotransmitter Transporters** • Acetylcholine • BGT-1 • Dopamine • GABA (GAT1, -2, -3) • VGAT • Glycine (Glyt 1, 2) • Glutamate (GLT1, GLAST, EAAC1) • Serotonin • Proline, Vesicular GABA Transporter (VGAT) • NET • Taurine • **Renal Physiology** Arginine Vasopressin Receptors 1/2 • Aquaporins (AQP 1-5) and Urea transporter (RUT2)

Western Blot recycling kit (Use the same blot to probe with multiple antibodies) **recycle blot at room temp in 5-10 min; No mercaptoethanol or heating required).**

NHE41-S-A-P 110519A

India Contact:

Life Technologies (India) Pvt. Ltd.

306, Aggarwal City Mall, Opposite M2K Pitampura, Delhi – 110034 (INDIA). Ph: +91-11-42208000, 42208111, 42208222, Mobile: +91-9810521400, Fax: +91-11-42208444

Email: customerservice@lifetechindia.com Website: www.lifetechindia.com