

Product Specification Sheet

NHERF2 Subunit Antibodies

Cat. # NHERF21-P	HumanNHERF2 control/blocking peptide # 1	SIZE: 100 ug
Cat. # NHERF21-A	Rabbit Anti-Human NHERF2 IgG # 1 (aff pure)	SIZE: 100 ug

Na⁺/ H⁺ exchanger regulatory factor (**NHERF1**) is a PDZ domain containing adaptor protein known to bind to various receptors, channels, cytoskeletal elements and cytoplasmic signaling proteins. A 358aa mainly expressed in kidney, liver and pancreas by NHERF gene (chr17). And is represented in another isoform, NHERF2. NHERF1 protein contains 2 tandem PDZ domains of approximately 90aa and a C-terminal sequence that binds several members of the ERM (ezrin-radixin-moesin) family of membrane cytoskeletal adaptors. The protein plays an important role in regulation of NHE3, turnover of G-protein coupled receptors, platelet derived growth factor receptor and ion transporters such as CFTR, Na⁺/Pi cotransporter, NaHCO₃ cotransporter and Trp channels.

NHERF1 overexpression in cancers and mutations in NHERF1 targets, Merlin, the product of tumor suppressor gene, also expressed in 2 isoforms, Merlin is involved in pathogenesis of benign tumors of the human nervous system, Merlin is named for its striking similarity with ERM family which thought to link cytoskeletal components with proteins in the cell membrane, It has a binding partner called Syntenin, which is an adapter protein that couples transmembrane proteoglycans to cytoskeletal components.

FUNCTION: Scaffold protein that connects plasma membrane proteins with members of the ezrin/moesin/radixin family and thereby helps to link them to the actin cytoskeleton and to regulate their surface expression. Necessary for cAMP-mediated phosphorylation and inhibition of SLC9A3. May also act as scaffold protein in the nucleus.

SUBCELLULAR LOCATION: Intracytoplasmic membrane; Peripheral membrane protein. Nucleus. Note=Nuclear, in a punctate pattern.

SIMILARITY: Contains 2 PDZ (DHR) domains.

Protein name Na(+)/H(+) exchange regulatory cofactor NHE-RF2

Synonyms NHERF-2

Tyrosine kinase activator protein 1

TKA-1

SRY-interacting protein 1

SIP-1

Solute carrier family 9 isoform A3 regulatory factor 2

NHE3 kinase A regulatory protein E3KARP

Sodium-hydrogen exchanger regulatory factor 2

Gene name Name: SLC9A3R2 Synonyms: NHERF2

Source of Antigen and Antibodies

Antigen	15-aa peptide from Human NHERF2/SLC9A3R2 ; protein accession #Q15599, refs 1 Designation (NHERF21-P, control peptide) conjugated to KLH; Epitope location~C-terminus
Ab Host/type	Rabbit, Polyclonal Aff pure IgG (cat # NHERF21-A) purified over the antigen column
2-ab	Cat # 20320, goat anti-rabbit IgG-HRP (AP, biotin, FITC conjugates also available)
-ve control	# 20009-1, Rabbit (non-immune) IgG, purified, suitable for ELISA, Western, IHC as -ve control

Form & Storage of Antibodies/Peptide Control

Affinity pure IgG

100 ug/100ul solution lyophilized powder

Supplied in Buffer: PBS+0.1% BSA

Reconstitute powder in PBS at 1mg/ml

Control/blocking peptide

100 ug/100 ul solution lyophilized powder

Supplied in Buffer: PBS pH 7.5,

Reconstitute powder in PBS at 1 mg/ml.

Storage

Short-term: unopened, undiluted liquid vials at -20OC and powder at 4oC or -20oC..

Long-term: at -20C or below in suitable aliquots after reconstitution. Do not freeze and thaw and store working, diluted solutions.

Stability: 6-12 months at -20oC or below.

Shipping: 4oC for solutions and room temp for powder

Recommended Usage

Western Blotting (1: 1-10 ug/ml for affinity pure antibody using ECL technique).

ELISA: Control peptide can be used to coat ELISA plates at 1 ug/ml and detected with antibodies (0.5-1 ug/ml for affinity pure).

Histochemistry & Immunofluorescence: Not tested. We recommend the use of aff pure IgG at 2-20 ug/ml.

Specificity & Cross-reactivity

The Human NHERF21-P control peptide is 100% identical in mouse, and rat. Antibody cross-reactivity in various species has not been studied. Control peptide, because of its low mol. Wt (<3 kDa), is not suitable for Western. It should be used for ELISA or antibody blocking experiments (use 5-10 ug control peptide per 1 ug of aff pure IgG or 1 ul antiserum) to confirm antibody specificity (see detailed protocol at: www.4adi.com/data/abblock.html).

General References: Yun, C.H (1998) JBC 273 (40), 25856-25863; Yun, C.H (1997) PNAS 94 (7), 3010-3015; Anita Murthy et al (1997) JBC 273, 1273-1276; Mehrdad Jannatipour et al (2001) JBC 276, 33093-33100; Edward J. Weinman et al (2001) Jour. Of Clinical Investigation, 108, 185-186.

*This product is for In vitro research use only.

Related material available from ADI

Antibodies and Peptides: NHERF isoforms, Merlin (Sch) isoforms, and Syntenin.

NHERF21-A-P

70829J

India Contact:

Life Technologies (India) Pvt. Ltd.

306, Aggarwal City Mall, Opposite M2K Pitampura, Delhi - 110034 (INDIA). Ph: +91-11-42208000, 42208111, 42208222, Mobile: +91-9810521400, Fax: +91-11-42208444

Email: customerservice@lifetechindia.com Website: www.lifetechindia.com