

Product Specification Sheet

Neuroigin 2 (NLGN2) Antibodies

Cat # NLGN21-P	Human NLGN2 Control/Blocking Peptide	SIZE: 100 µg
Cat # NLGN21-A	Rabbit anti-human NLGN2 IgG (affinity pure)	SIZE: 100 µg

Neuroigins are Type I membrane proteins enriched in synaptic plasma membranes and clustered in synaptic clefts and postsynaptic densities. They have been characterized as neuronal cell surface proteins and are thought to be involved in cell-cell-interactions by forming intercellular junctions through binding to beta-neurexins. They play a major role in the formation or maintenance of synaptic junctions. They are also thought to be involved in the specification of excitatory synapses. Neuroigins interact with **beta-neurexins** and this interaction is involved in the formation of functional synapses.

NLGN2 (rat: 836aa; mouse: 836aa; human 835-aa, 91kDa, human chromosome 17p13). Neuroigin 2 is exclusively localized to inhibitory synapses. It interacts with neurexin 1-beta, neurexin 2-beta and neurexin 3-beta.

Source of Antigen, Antibodies

Antigen	18- aa peptide of Human NLGN2 (Protein accession # Q8NFZ4 ; ref. 1); designated as NLGN21-P control/blocking peptide conjugated to KLH; epitope location ~C-terminus, cytoplasmic
Antibody host/type	Rabbit, Polyclonal IgG (Cat # NLGN21-A), purified over antigen-Agarose
Secondary Ab	Cat # 20320, goat anti-rabbit IgG-HRP (AP, biotin, FITC conjugates also available).
Negative Control Ab	Non-immune rabbit IgG (Cat # 20009-1) to be used as -ve control for ELISA, WB, IHC etc.

Form & Storage of Antibodies/Peptide Control

Affinity pure IgG

100 ug/100ul solution lyophilized powder
Supplied in **Buffer:** PBS+0.1% BSA
Reconstitute powder in PBS at 1mg/ml

Control/blocking peptide

100 ug/100 ul solution lyophilized powder
Supplied in Buffer: PBS pH 7.5,
Reconstitute powder in PBS at 1 mg/ml.

Storage

Short-term: unopened, undiluted vials for less than a week at 4°C.

Long-term: at -20°C or below in suitable aliquots after reconstitution. Do not freeze and thaw and store working, diluted solutions.

Stability: 6-12 months at -20°C or below.

Shipping: 4°C for solutions and room temp for powder.

Recommended Usage

Western Blotting: 1-10 µg/ml; using affinity pure antibody (chemiluminescence technique).

ELISA: 1:100K; using 50-100 ng control peptide/well.

Histochemistry & Immunofluorescence: Not tested; we recommend the use of affinity purified antibody at 2-10 µg/ml.

Specificity & Cross-reactivity

Human NLGN21-P peptide sequence is 100% conserved in both mouse and rat NLGN2 protein. Antibody cross-reactivity in various species is not known. The control peptide, because of its low mol. Wt (<3 kDa), is not suitable for Western. It should be used for ELISA or antibody blocking experiments (use 5-10 ug control peptide per 1 ug of aff pure IgG or 1 ul antiserum) to confirm antibody specificity (see detailed protocol at the web site).

General References:

(1) Nagase T., et al., (2000) DNA Res. 7: 65-73

List of related items, data sheets, and publications, using ADI antibodies is posted on the web site

*This product is for in vitro research use only.

Related material available from ADI

Antibodies to NLGN1, NLGN3 and NLGN4

NLGN21-A-P 80115J

India Contact:

Life Technologies (India) Pvt. Ltd.

306, Aggarwal City Mall, Opposite M2K Pitampura, Delhi - 110034 (INDIA). Ph: +91-11-42208000, 42208111, 42208222, Mobile: +91-9810521400, Fax: +91-11-42208444
Email: customerservice@lifetechindia.com Website: www.lifetechindia.com