

Product Specification Sheet

NPT2a Antibodies

Cat # NPT28-P	Human NPT2a control/blocking peptide # 8	SIZE: 100 ug
Cat # NPT28-S	Chicken Anti-Human NPT2a antiserum #8	SIZE: 100 ul

Inorganic phosphate (Pi) levels are hormonally regulated that affects the physiological activity of bone, kidney, and small intestine. Majority of the Pi is absorbed in the small intestine and reabsorbed in the proximal tubules in the kidney. At least 4 groups of structurally and functionally related proteins are involved in Pi transport: **Type I-related** NaPi transporters designated **NPT1**, **Npt1**, and **NaPi-1** respectively in humans, mouse, and rabbit are expressed in the kidney and liver. Its expression and activity are not regulated by Pi deprivation or parathyroid hormone (PTH) and its role in Pi-homeostasis is not clear. **Type IIa-related** cotransporters, designated **NaPi-2** in rat, **NaPi-3** or **NPT2** in humans, **NaPi-4** in opossum, **NaPi-5** in flounder vessel, **NaPi-6** or **Npt-2** in mouse, and **NaPi-7** in rabbit, is the primary target for Pi regulation by dietary, hormonal, and tubular Pi reabsorption. Deletion of Npt2 gene produces severe Pi wasting. Type II transporters are expressed in kidney, brain, lung, bone and small intestine. **Type IIb**, designated as **NaPi-IIB** or **NaPi-2b** in rat/mouse, and **NaPi-3b** in human, is closely related isoform of the NaPi-2 family. It is expressed in small intestine and lung. **Type III NaPi** transporters, originally described as a family of cell surface receptors for gibbon ape leukemia virus (**GALV**) and murine amphotropic retrovirus (**A-MuLV**), share very low (<20%) sequence homology with Type I and II proteins, and are found in most tissues. Human homologs are now designated as **Pit1** (also known as GALV receptors or GLVR1) and **Pit2** (**A-MuLV** receptors or GLVR2 or Ram1). Pit1 and Pit2 are ~62% related and predicted to contain 10 TM domains that is in contrast with Type 1/2 transporters (6-8 TM). The N and C-termini are predicted to be cytoplasmic.

Source of Antigen and Antibodies

Antigen	Human Type-IIa, NPT2/Napi-3 is ~640 aa (chromosome 8) and mouse and rat 637 aa proteins. A 21 AA synthetic peptide (designated NPT28-P; control peptide) within the putative, cytoplasmic, C-terminus of Human NPT2a was synthesized, conjugated to KLH
Ab Host/type	Rabbit, Polyclonal antiserum # HO22-S and IgG, purified over antigen-agarose (Cat # HO22-A). An antibody to this antigenic peptide is also produced in rabbits (Cat # NPT27-S & NPT27-A).
2-Ab	Cat # 20320, goat anti-rabbit IgG-HRP (AP, biotin, FITC conjugates also available).
-ve control IgG	Cat # 20009-1, Rabbit (non-immune) Serum IgG, purified, suitable for ELISA, Western, IHC as -ve control

Form & Storage of Antibodies/Peptide Control

Antiserum (unpurified)
100ul solution lyophilized powder
Supplied 0.05% azide, **Reconstitute** powder in 100 ul PBS

Affinity pure IgG

100 ug/100ul solution lyophilized powder
Supplied in **Buffer:** PBS+0.1% BSA
Reconstitute powder in PBS at 1mg/ml

Control/blocking peptide

100 ug/100 ul solution lyophilized powder
Supplied in Buffer: PBS pH 7.5,
Reconstitute powder in PBS at 1 mg/ml.

Storage

Short-term: unopened, undiluted liquid vials at -20OC and powder at 4oC or -20oC..

Long-term: at -20C or below in suitable aliquots after reconstitution. Do not freeze and thaw and store working, diluted solutions.

Stability: 6-12 months at -20oC or below.

Shipping: 4oC for solutions and room temp for powder

Recommended Usage

Western Blotting 1:1K-5K for antiserum using Chemiluminescence technique. NPT2a apparent mol wt is 80-90 kDa, and under reducing conditions 2 bands of 40-50 kDa are observed. It may also form multimeric structures (170 and 200 kDa). (see Murer et al 2000)

ELISA: Control peptide can be used to coat ELISA plates at 1 ug/ml and detected with antibodies (1:10-50K for neat serum).

Histochemistry & Immunofluorescence: not tested.

Specificity & Cross-reactivity

The NPT28-P peptide is 100% conserved in human, rat, mouse, rabbit, and 86% in ovine, 81% bovine, and 71% in opossum NPT2a. No significant sequence homology of NPT25-P is observed with NPT2b or other Pi-transporters (Type-I, III, or IV) transporters. Antibody crossreactivity in various species or isoform is not established. Control peptide, because of its low mol. Wt (<3 kDa), is not suitable for Western. It should be used for ELISA or antibody blocking experiments (use 5-10 ug control peptide per 1 ug of aff pure IgG or 1 ul antiserum) to confirm antibody specificity (see detailed protocol see detailed protocol at the web site).

General References:

- (1). Magnin S et al (1993) PNAS 90, 5979; Collins JF et al (1996) FASEB J 8, 862; Hartman CM et al (1996) PNAS 93, 7409.
- (2) Murer H et al (2000) Physiol. Rev. 80, 1373 (review); Werner A et al (2001) Am. J. Physiol. Integ. Comp. Physiol. 280, R301

**This product is for In vitro research use only.*

Related material available from ADI

Anti-Napi I-III, Sialin, EAAC1, GLT1, EAAT4, EAAT5, GTRAP41, NPT25, VGLUT1/BNPI, VGLUT-2/DNPI & GABA Transporters
NPT28-S-P 71213A

India Contact:

Life Technologies (India) Pvt. Ltd.

306, Aggarwal City Mall, Opposite M2K Pitampura, Delhi - 110034 (INDIA). Ph: +91-11-42208000, 42208111, 42208222, Mobile: +91-9810521400, Fax: +91-11-42208444

Email: customerservice@lifetechindia.com Website: www.lifetechindia.com