

Product Specification Sheet

Neurotensin Receptor 1 (NTR1) Antibodies

| | | |
|---------------------|---|---------------------|
| Cat. NTR11-S | Rabbit Anti-Mouse NTR1 antiserum (Ab # 1) | SIZE: 100 ul |
| Cat. NTR11-A | Rabbit Anti-Mouse NTR1 Ig G # 1 (affinity pure) | SIZE: 100 ug |
| Cat. NTR11-P | Mouse NTR1 Control/blocking peptide # 1 | SIZE: 100 ug |

Neurotensin (NT) is an endogenous tridecapeptide neurotransmitter that triggers hypothermic and naloxone-insensitive analgesic responses, whereas peripheral effects include hypotension, decrease gastric acid release, potentiation of lipid digestion. NT causes contraction of smooth muscle. NT is widely distributed throughout the CNS. NT gene (human chromosome 12q21) produces both NT and neuromedin N (NN). NT initiates its biological action by interacting with two distinct G-protein coupled receptors (**NTR1 and NTR2**). Recently, a third receptor **NTR3** has been identified that is identical to **gp95/sortilin** and it is not coupled via the G-proteins. All three receptors bind NT through its C-terminal hexapeptide sequence (**8 RRPYL 13**). Biologically active **NT (NT8-13)** has also been shown to interact with the extracellular domain 3 (between TM6-7) of NTR1.

NTR1/NTRH/NTSR1/NT1 (mouse/rat 424 aa; human 418 aa, chromosome 20q1; ~84% interspecies sequence homology) is the levocabastine-insensitive, high affinity receptor for NT. It has the typical structure of GPCR: 7 TM domains with extracellular N-terminus and cytoplasmic C-terminus. It is shown to mediate a number of peripheral and central NT responses, including the neuroleptic-like effects of the peptide. It is highly expressed in the brain, small intestine, and other peripheral tissues.

FUNCTION: Catalyzes the phosphorylation of the 3'-hydroxyl group of dephosphocoenzyme A to form coenzyme A (By similarity).

SUBCELLULAR LOCATION: Cytoplasm (By similarity).

SIMILARITY: Belongs to the coaE family.

Protein name Dephospho-CoA kinase

Synonyms EC 2.7.1.24, Dephosphocoenzyme A kinase

Gene name Name: coaE

OrderedLocusNames: TP_0296

Source of Antigen and Antibodies

| | |
|---------------------|--|
| Antigen | 18aa peptide of Mouse NTR1; (protein accession #O88319, refs 1) Designated (NTR11-P or control peptide) , conjugated to KLH; Epitope location ~ Mid-region, Extracellular |
| Ab Host/type | Rabbit, Polyclonal Aff pure (cat # NTR11-A) purified over the antigen column |
| Ab Format | Cat # 20320, goat anti-rabbit IgG-HRP (AP, biotin, FITC conjugates also available |
| -ve control | # 20009-1, Rabbit (non-immune) IgG, purified, suitable for ELISA, Western, IHC as -ve control |

Form & Storage of Antibodies/Peptide Control

Antiserum (unpurified)

100ul solution lyophilized powder
Supplied in Buffer: 0.05% azide
Reconstitute powder in 100 ul PBS

Affinity pure IgG

100 ug/100ul solution lyophilized powder
Supplied in **Buffer:** PBS+0.1% BSA

Reconstitute powder in PBS at 1mg/ml

Control/blocking peptide

100 ug/100 ul solution lyophilized powder
Supplied in Buffer: PBS pH 7.5,

Reconstitute powder in PBS at 1 mg/ml.

Storage

Short-term: unopened, undiluted liquid vials at 20°C and powder at 4°C or -20°C..

Long-term: at -20°C or below in suitable aliquots after reconstitution. Do not freeze and thaw and store working, diluted solutions.

Stability: 6-12 months at -20°C or below.

Shipping: 4°C for solutions and room temp for powder

Recommended Usage

Western Blotting (1:1K-5K for neat serum and 1-10 ug/ml for affinity pure using Chemiluminescence technique). NTR1 has been shown to be 52-54 kDa in rat brain (1). NTR1 may be differentially glycosylated.

ELISA (1:100K; using 50-100 ng control peptide/well).

Histochemistry & Immunofluorescence: Not tested. we recommend the use of affinity purified antibody at 2-20 ug/ml in formaldehyde fixed tissue.

Specificity & Cross-reactivity

The mouse NTR11-P sequence 84% conserved in rat and human NTR1. No significant sequence homology exist with other NTR2 or NTR3 or other GPCR. Antibody cross-reactivity in various species has not been studied. Control peptide, because of its low mol. Wt (<3 kDa), is not suitable for Western. It should be used for ELISA or antibody blocking experiments (use 5-10 ug control peptide per 1 ug of aff pure IgG or 1 ul antiserum) to confirm antibody specificity

General References: (1) Vita N et al (1993) FEBS Lett. 317, 139-142; Tanaka K et al (1990) Neuron 4, 847-854; Labbe-Julie C et al (1994) J Pharmacol. Exp. Ther. 268, 328; Boudin H et al (2000) J Compo. Neurol. 425, 45-57.

*This product is for in vitro research use only.

Some New Antibodies from ADI...

- Neuromedin U, NMUR1 and NMUR2 antibodies, Neurotensin receptors NTR1-3, Ghrelin receptors, Orexin and orexin receptors, CART, and Leptin receptors

Western Blot recycling kit (Use the same blot to probe with multiple antibodies) **recycle blot in 5-10 min.**

Study distribution of proteins in **pre-made protein blots of brain, kidney, GI-tract, and major tissues**

NTR11-S-A-P

70907J

India Contact:

Life Technologies (India) Pvt. Ltd.

306, Aggarwal City Mall, Opposite M2K Pitampura, Delhi - 110034 (INDIA). Ph: +91-11-42208000, 42208111, 42208222, Mobile: +91-9810521400, Fax: +91-11-42208444

Email: customerservice@lifetechindia.com Website: www.lifetechindia.com