

Product Specification Sheet

**Organic Anion Transporter 3 (OAT3) Antibodies**

Cat. # OAT31-P	Rat OAT3 control/blocking Peptide	<b>SIZE:</b> 100 ug
Cat. # OAT31-S	Rabbit Anti-rat OAT3 antiserum	<b>SIZE:</b> 100 ul
Cat. # OAT31-A	Rabbit Anti-rat OAT3 IgG (aff pure)	<b>SIZE:</b> 100 ug

Mammalian kidney and liver are critical in maintaining physiological ionic environment. Kidney specializes in removing toxins, drugs, and other organic anions from the blood by a process called "renal secretion". Besides kidney, anionic substrates are also transported in other organs, e.g., choroid plexus, eye, airway, and placenta. Several multispecific **OATs** (OAT1-3, OAT-K1 and OATK2) and **OATPs** (organic anion transporting polypeptides; **oatp1-3**), have been cloned and characterized from various tissues. OATPs family of proteins are very similar in sequence and secondary protein structure (up to 12 transmembrane domains with cytoplasmic N and C-terminus).

Recently **OAT3** (rat 536-aa, human 568 aa) has been cloned that is most closely related to OAT1 (~49% identity). Rat OAT3 is expressed in the liver, brain, kidney and eye. OAT3 also mediates Na-independent uptake of several organic anions (PAH, ochratoxin, and estrone sulfate, etc). OAT3 shows 92% homology with mouse **Roct** (reduced in osteosclerosis transporter), possibly an ortholog of rat OAT3. As compared to OAT1-2, strong expression of OAT3 in brain suggests its role in removing organic anion in the brain.

**Source of Antigen and Antibodies**

<b>Antigen</b>	16aa peptide of Rat OAT3 ; (gene accession # Np112622.1) <b>Designated (OAT31-P or control peptide)</b> conjugated to KLH; Epitope location ~N-terminus
<b>Ab Host/type</b>	Rabbit, polyclonal Unpurified antiserum (cat #OAT31-S) Aff pure IgG (cat #OAT31-A)
<b>2-ab</b>	Goat Anti-rabbit IgG-HRP cat # 20320 (AP, biotin, FITC conjugates also available)
<b>-ve control</b>	# 20009-1, Rabbit (non-immune) IgG, purified, suitable for ELISA, Western, IHC as -ve control

**Form & Storage of Antibodies/Peptide Control**

**Antiserum (unpurified)**

100ul solution lyophilized powder  
Supplied in Buffer: 0.05% azide  
**Reconstitute** powder in 100 ul PBS

**Affinity pure IgG**

100 ug/100ul solution lyophilized powder  
Supplied in **Buffer:** PBS+0.1% BSA  
**Reconstitute powder** in PBS at 1mg/ml

**Control/blocking peptide**

100 ug/100 ul solution lyophilized powder  
Supplied in Buffer: PBS pH 7.5,  
**Reconstitute powder** in PBS at 1 mg/ml.

**Storage**

**Short-term:** unopened, undiluted liquid vials at 20°C and powder at 4°C or -20°C..

**Long-term:** at -20°C or below in suitable aliquots after reconstitution. Do not freeze and thaw and store working, diluted solutions.

**Stability:** 6-12 months at -20°C or below.

**Shipping:** 4°C for solutions and room temp for powder

**Recommended Usage**

**Western Blotting** (1:1K-5K for neat serum and 1-10 ug/ml for affinity pure antibody using ECL technique).

**ELISA:** Control peptide can be used to coat ELISA plates at 1 ug/ml and detected with antibodies (1:10-50K for neat serum and 0.5-1 ug/ml for affinity pure).

**Histochemistry & Immunofluorescence:** We recommend the use of affinity purified antibody at 2-20 ug/ml. See refs 2.

**Specificity & Cross-reactivity**

The 16 AA rat OAT31-P control peptide is 93% conserved in mouse, 87% in, rabbit and human OAT3. No significant sequence homology is detected with other OATs or other proteins. Antibody cross-reactivity in various species has not been studied. Control peptide, because of its low mol. Wt (<3 kDa), is not suitable for Western. It should be used for ELISA or antibody blocking experiments (use 5-10 ug control peptide per 1 ug of aff pure IgG or 1 ul antiserum) to confirm antibody specificity

**General References:** Kushuhara h et al (1999) J Biol. Chem. 274, 13675-13680; Race JE et al (1999) BBRC 255, 508

(2) Citations of ADI's Antibodies (see web site for updated list)

OAT3 Bahn A 2005 Am J Physiol Cell Physiol, 289: C1075 - C1084 IF

\*This product is for In vitro research use only.

**Related material available from ADI**

Antibodies OAT1-7 and CLC-K1; KCCL1-3; AQP-9 and RUT; OCT and OAT, AE-3, and NACX

Study distribution of proteins in pre-made **Kidney blots** from 7 defined regions of rat kidney

**Recycle your blot in Just 5-10 min. (use the same strip for various OATs)**

OAT31-S-A-P

60908A

**India Contact:**

**Life Technologies (India) Pvt. Ltd.**

306, Aggarwal City Mall, Opposite M2K Pitampura, Delhi - 110034 (INDIA). Ph: +91-11-42208000, 42208111, 42208222, Mobile: +91-9810521400, Fax: +91-11-42208444  
Email: [customerservice@lifetechindia.com](mailto:customerservice@lifetechindia.com) Website: [www.lifetechindia.com](http://www.lifetechindia.com)