

OREXIN-2 Receptor (Hypocretin-2 Receptor) Antibodies

Cat. OX2R22-S	Rabbit Anti-Hhuman Orexin-2 Receptor Antiserum #2	SIZE: 100 ul
Cat. OX2R22-A	Rabbit Anti-Human Orexin-2 Receptor IgG #2 (aff pure)	SIZE: 100 ug
Cat. OX2R22-P	Human Orexin-2 Receptor Control peptide #2	SIZE: 100 ug

Several peptides associated with feeding behavior have been reported recently. Orexins (**Orexin-A** and **Orexin-B**) are a family of hypothalamic neuropeptides selectively expressed in the hypothalamus (1-2). Orexin-A and Orexin-B are derived from the same precursor (Prepro-orexin) by proteolytic cleavage. **Prepro-orexin** is 130 amino acid long peptide with a putative 33-AA secretory sequence, a hydrophobic core followed by residues with small polar side chains. The expression was detected in brain and to a small extent in testis (1-2). These neuropeptides bind and activate two closely related **Orexin receptors**--G-protein coupled receptors (GPCRs) **OX1R** and **OX2R**. Rat and human OX2R are 460 aa, and 444 aa, respectively (1-2)

Source of Antigen and Antibodies

Antigen	19-aa peptide from human OX2R/hcrtr2 (gene accession # O43614 , refs 1); Designation (#OX2R22-P , control/blocking peptide) conjugated to KLH
Epitope Location	~N-terminus Extracellular domain
Ab Host/type	Rabbit, Polyclonal unpurified antiserum (#OX2R22-S) and IgG, purified over antigen-agarose (Cat # OX2R22-A)
2-Ab	Cat # 20320, goat anti-rabbit IgG-HRP (AP, biotin, FITC conjugates also available).
-ve control	# 20009-1, Rabbit (non-immune) IgG, purified, suitable for ELISA, Western, IHC as -ve control

Form & Storage of Antibodies/Peptide Control

Antiserum (unpurified)

100ul solution lyophilized powder
Supplied in Buffer: 0.05% azide
Reconstitute powder in 100 ul PBS

Affinity pure IgG

100 ug/100ul solution lyophilized powder
Supplied in Buffer: PBS+0.1% BSA
Reconstitute powder in PBS at 1mg/ml

Control/blocking peptide

100 ug/100 ul solution lyophilized powder
Supplied in Buffer: PBS pH 7.5,
Reconstitute powder in PBS at 1 mg/ml.

Storage

Short-term: unopened, undiluted liquid vials at -200C and powder at 4oC or -20oC..

Long-term: at -20C or below in suitable aliquots after reconstitution. Do not freeze and thaw and store working, diluted solutions.

Stability: 6-12 months at -20oC or below.

Shipping: 4oC for solutions and room temp for powder

Recommended Usage

Western Blotting (1:1K-5K for neat serum and 1-10 ug/ml for affinity pure using ECL). (see published refs using this antibody in 2).

ELISA (1:10K-1:100K; using 50-100 ng of control peptide/well).

Histochemistry: We recommend the use of affinity-purified antibody at 2-20 ug/ml (see published refs. 2)

Specificity & Cross-reactivity

Human OX2R22-P peptide is 94% in dog and pig, 84% in rat and 78% in mouse OX2R. No significant homology is seen with OX1R or other G-protein-coupled receptors. Antibody reactivity in all species is not established. Control peptide, because of its low mol. Wt (<3 kDa), is not suitable for Western. It should be used for ELISA or antibody blocking experiments (use 5-10 ug control peptide per 1 ug of aff pure IgG or 1 ul antiserum) to confirm antibody specificity (see detailed protocol at: www.4adi.com/data/abblock.html).

General References: (1) Sakurai, T. et al. (1998) *Cell*, **92**, 573-585; DeLecca L et al (1998) *PNAS* **95**, 322-327

(2) Citations of ADI's Antibodies (see web site for updated list)

Beiras-Fernández A, 2004, *J Anatomy* **204**, 117-122, IHC,
Naslun E, 2002, *AJP Gastrointest Liver Physiol* **282**: G470-G479
Blanco, M., 2001, *J. Clin. Endocrinol. Metab.* **86**: 3444
Blanco, M., 2001, *Regulatory Peptide* in press
Beiras-Fernández A, 2004, *Journal of Anat.* **204**, 117
Caillol M 2003 *Brain Res.* **960**, 48-61
Lopez M, 1999, *Endocrinol.* **140**, 5991-5994
Blanco M, 2001, *J Clin. Endocrinol. Metabol.* **86**, 1616
Kirchgessner, K., 1999, *Neuron* **24**: 941-951

*This product is for *in vitro* research use only.

Related material available from ADI:

OX2R22-S-A-P 70622A