

Product Specification Sheet

**Mouse Obestatin Antibodies**

|                     |   |                     |
|---------------------|---|---------------------|
| <b>Cat. OXA11-S</b> | Rabbit Anti-Mouse/Rat Orexin-A antiserum        | <b>SIZE:</b> 100 ul |
| <b>Cat. OXA11-A</b> | Rabbit Anti-Mouse/Rat Orexin-A, Ig G (aff pure) | <b>SIZE:</b> 100 ug |
| <b>Cat. OXA11-P</b> | Mouse/Rat Orexin-A Control peptide              | <b>SIZE:</b> 100 ug |

Several peptides associated with feeding behavior have been reported recently. Orexins (**Orexin-A** and **Orexin-B**) are a family of hypothalamic neuropeptides selectively expressed in the hypothalamus (1-2). Orexin-A and Orexin-B are derived from the same precursor (Prepro-orexin) by proteolytic cleavage. **Prepro-orexin** is 130 amino acid long peptide with a putative 33 AA secretory sequence, a hydrophobic core followed by residues with small polar side chains. The expression was detected in brain and to a small extent in testis (1-2). These neuropeptide bind and activate two closely related **Orexin receptors**--G-protein coupled receptors (GPCRs) **OX1R** and **OX2R**.

**Source of Antigen and Antibodies**

|                        |  |
|------------------------|--|
| <b>Antigen</b>         | 18-aa peptide of mouse Orexin-A (gene accession # 055232) ( <b>designated as OXA11-P, control peptide</b> ) conjugated to KLH. The peptide has two disulfides (Cys 6-Cys 12 & Cys 7-Cys 14). ; epitope location ~ N-terminus |
| <b>Ab Host/type</b>    | Rabbit Polyclonal, unpurified antiserum (#OXA11-S) and IgG, purified over antigen-agarose (Cat # PXA11-A) supplied in PBS+0.1% BSA+0.05% azide   |
| <b>2-Ab</b>            | Cat # 20320, goat anti-rabbit IgG-HRP (AP, biotin, FITC conjugates also available).  |
| <b>-ve control IgG</b> | # 20009-1, Rabbit (non-immune) IgG, purified, suitable for ELISA, Western, IHC as -ve control  |

**Form & Storage of Antibodies/Peptide Control**

**Antiserum (unpurified)**

100ul solution lyophilized powder  
Supplied in Buffer: 0.05% azide  
**Reconstitute powder in 100 ul PBS**

**Affinity pure IgG**

100 ug/100ul solution lyophilized powder  
Supplied in **Buffer:** PBS+0.1% BSA  
**Reconstitute powder in PBS at 1mg/ml**

**Control/blocking peptide**

100 ug/100 ul solution lyophilized powder  
Supplied in Buffer: PBS pH 7.5,  
**Reconstitute powder in PBS at 1 mg/ml.**

**Storage**

**Short-term:** unopened, undiluted liquid vials at -200C and powder at 4oC or -20oC..

**Long-term:** at -20C or below in suitable aliquots after reconstitution. Do not freeze and thaw and store working, diluted solutions.

**Stability:** 6-12 months at -20oC or below.

**Shipping:** 4oC for solutions and room temp for powder

**Recommended Usage**

**Western Blotting** (1:1K-5K for neat serum and 1-10 ug/ml for aff pure using Chemiluminescence technique).

**ELISA** (1:10K-1:100K; using 50-100 ng of control peptide/well).

**Histochemistry & Immunofluorescence:** We recommend the use of affinity purified antibody at 2-20 ug/ml (see published refs 2)

**Specificity & Cross-reactivity**

Mouse OXA11-P control peptide is 100% identical in rat, pig, ovine, bovine, dog, human and 81% in chicken. Anti-OXA11 reacts with the full-length (1-33 aa) Orexin-A. The OXA11-P epitope is not found in Orexin-B. Since Orexin-A and Orexin-B are derived from an unprocessed pre-proorexin (130 aa), the anti-Orexin-A antibodies may also react with pre-pro orexin. Antibodies made to prepro-orexin A are also available. Antibody crossreactivity in various species is not established. Control peptide, because of its low mol. Wt (<3 kDa), is not suitable for Western. It should be used for ELISA or antibody blocking experiments (use 5-10 ug control peptide per 1 ug of aff pure IgG or 1 ul antiserum) to confirm antibody specificity (see detailed protocol see detailed protocol at the web site)..

**General References: (1)** Sakurai, T. et al. (1998) *Cell*, **92**, 573-585; DeLecca L et al (1998) *PNAS* 95, 322-327

**(2) Citations of ADI's Antibodies** (see web site for updated list)

Ciriello J, 2003, *Brain Research*, 967, Pages 123-131; Ouedraogo R., 2003, *Diabetes*, Jan 2003; 52: 111- 117; deOliveira CVR, 2003, *Am J Physiol Heart Circ Physiol* 284: H1369; Satoh S, 2003, *Eur. J. Neuroscience* 17, 1635; Davis SF, 2003, *J. Neurosci.*, 23: 3844 – 3854; Ciriello J, 2003, *Brain Research*, 991, 84-95; Ehrstrom M., 2003, *Am J Physiol Gastrointest Liver Physiol*, 285, 688; Ciriello J, 2003, *Am J Physiol Regul. Int. Comp Physiol*, 284: 1611; deOliveira CVR, 2003, *Brain Research* 978, 14-23

\*This product is for *in vitro* research use only.

OXA11-S-A-P 100811A

Alpha Diagnostic Intl Inc., 6203 Woodlake Center Dr, San Antonio, TX 78244, USA; **Email:** [service@4adi.com](mailto:service@4adi.com)

**India Contact:**

**Life Technologies (India) Pvt. Ltd.**

306, Aggarwal City Mall, Opposite M2K Pitampura, Delhi – 110034 (INDIA). Ph: +91-11-42208000, 42208111, 42208222, Mobile: +91-9810521400, Fax: +91-11-42208444  
Email: [customerservice@lifetechindia.com](mailto:customerservice@lifetechindia.com) Website: [www.lifetechindia.com](http://www.lifetechindia.com)