


Product Specification Sheet

Porcine Circovirus 2 replicase protein (PCV2-rep/ORF1) Antibodies and Controls

| | | |
|-------------------|---|--------------------|
| - Cat # PCV2R11-S | Rabbit Anti-Porcine Circovirus 2 replicase protein (PCV2-rep/ORF1) antiserum | SIZE:100 ul |
| - Cat # PCV2R11-C | Recombinant Porcine Circovirus 2 replicase protein (PCV2-rep) control for western blot | SIZE:100 ul |

The viral genus **Circovirus** is a part of the family of Circoviridae and contains viral species with non-enveloped, circular ssDNA genomes. These viruses are the smallest viruses replicating autonomously in nucleus of host cells of vertebrate species, including pigs, dogs, and birds such as pigeons, and ducks. The virions of Circoviruses are surprisingly small, with diameters ranging from 17 up to 22 nm. **Porcine circovirus (PCV)** is a small, non-enveloped virus (17 nm in diameter) with a single stranded circular DNA genome. It causes worldwide infection in swine and is highly contagious. Porcine circovirus belong to the genus Circovirus of the Circoviridae family. The virus targets the lymphoid tissue and causes immunosuppression in the host. Pigs affected may experience increased mortality, poor growth, and weight loss, progressing to the level of severe thinning and weakness between 5 to 14 weeks of age. Two species of PCV have been characterized, PCV1 and PCV2. Although PCV1 was initially described as an infant of porcine kidney cell line (PK15), it shows no pathogenic traits. Recently, PCV2 has been identified as being associated with a new condition in pigs, the post weaning multi-systemic wasting syndrome (PMWS), first described in Western Canada. This syndrome is characterized clinically by progressive weight loss, dyspnea, and jaundice and pathologically by lymphadenopathy, interstitial pneumonia, hepatitis, and nephritis. Similar syndromes have also been recently reported in the United States, Europe, and Asia. Other diseases, such as porcine dermatitis and nephropathy syndrome (PDNS), proliferative and necrotizing pneumonia (PNP), perina-talmiocarditis and reproductive failures are also associated with PCV2. PMWS is commonly diagnosed on the continents of North America, Europe and Asia.



The PCV genome (~1.76 kb) has three main **open reading frames (ORFs)** that encode well characterized proteins: ORF1 encodes the viral **replicase**

proteins (REP and REP') that are essential for rolling circle replication. The replicases differ in that Rep is the full ORF1 transcript of 312-aa, whereas Rep' is a truncated form of ORF1 as a result of splicing and is only 168-aa in length. The two replicase enzymes that are created from ORF1, Rep and Rep', are conserved between the two types of PCV, and are part of the early phase of the virus.

ORF2 encodes the viral capsid protein (Cap) involved in host immune response. ORF2 differ slightly between PCV-1 and PCV-2. Antibodies to ORF2 are found in experimental PCV2 infection. ORF3 encodes an apoptotic protein, which is essential for the development of the viral pathogenesis. The non-pathogenic PCV1 and the pathogenic PCV2 share ~68-76% protein identity (~90% DNA). PCV2 are divided into five genotypes (PCV2a, PCV2b, PCV2c, PCV2d, and PCV2e) that are ~95% conserved. PCV2a was long the predominant strain in the global pig population and is also the basis of all major commercially available PCV2 vaccines. Currently, PCV2b is the most commonly identified strain in pigs. Diagnosis of PCV is confirmed by finding the typical histological lesions and demonstration of viral antigen or DNA within the lesions by immunohistochemical (IHC) or in situ hybridization methods (ISH) or by ELISA.

Source of Antigen and Antibodies

| | |
|------------------------|---|
| Antigen | Recombinant purified PCV2 Rep protein (cat# PCV2R15-R-10) |
| Ab Host/type | Rabbit, Polyclonal antiserum (Cat# PCV2R11-C) supplied in 0.05% azide |
| 2-Ab | Goat Anti-rabbit IgG-HRP cat # 20320 (AP, biotin, FITC conjugates) |
| -ve control IgG | # 20009-1 , Rabbit (non-immune) IgG, purified, suitable for ELISA, Western, IHC as -ve control |

Cat# PCV2R11-S, positive control

PCV2 Rep protein was expressed in E. coli as his-tag fusion protein (full length, >95%, ~36 KDa). Purified PCV2 Rep protein for Western blot +ve control (Cat# **PCV2R11-C**) is supplied in SDS-PAGE sample buffer (reduced). Load 10 ul/lane of # **PCV2R11-C** for good visibility with antibody

Cat# PCV2R11-S. Store at -20°C in suitable size aliquots. SDS may crystallize in cold conditions. It should redissolve by warming before taking it from the stock. It should be heated once prior to loading on gels. If the product has been stored for several weeks, then it may be preferable to add 5 ul of fresh 2x sample buffer per 10 ul of the # **PCV2R11-C** solution prior to heating and loading on gels. This preparation is not biologically active. It is not suitable for ELISA or other applications where native protein is required. Do not freeze, thaw, or heat repeatedly

Form & Storage of Antibodies/Peptide Control

Antiserum

100 ul solution lyophilized powder

Buffer: PBS+0.05% azide

Reconstitute powder 100 ul of PBS.

Storage

Short-term: unopened, undiluted vials for less than a week at 4°C.

Long-term: at -20°C or below in suitable aliquots after reconstitution. Do not freeze and thaw and store working, diluted solutions.

Stability: 6-12 months at -20°C or below.

Shipping: 4°C for solutions and room temp for powder.

Recommended Usage

Western Blotting: An initial dilution of 1:500-2K is recommended for Western. Purified PCV2Rep is ~36 kda. Users must optimize antibody dilution depending upon the nature of samples and other technical conditions.

ELISA (1:10-50K; using 50-100 ng antigen/well).

Histochemistry & Immunofluorescence: not tested.

Specificity & Cross-reactivity

This antibody reacts with PCV2 replicase protein and recombinant protein. Cross reactivity with other proteins has not been established. Antibodies and recombinant proteins to PCV2 ORF2 and replicase protein are available for control studies.

References: Shuai J (2007) Virus Genes 35, 619-627; Seo HW (2014) Vet. J. 200, 65-70. Ellis J (2014) Veterinary Pathology 51 (2): 315-327; G. Misinzo, et al. (2005). J Gen Virol. 86 (7): 2057-2068.

*This product is for In vitro research use only.

Related material available from ADI

| | |
|--------------|---|
| PCV2C21-C | Recombinant Porcine Circovirus 2 Capsid protein (PCV2-ORF2) control for western blot |
| PCV2C21-S | Rabbit Anti-Porcine Circovirus 2 Capsid protein (PCV2-ORF2) antiserum |
| PCV2C25-R-10 | Recombinant (E.coli) Porcine Circovirus 2 capsid (PCV2-ORF2) protein (his tag, 26 kDa) purified |
| PCV2R11-C | Recombinant Porcine Circovirus 2 replicase protein (PCV2-rep/ORF1) control for western blot |
| PCV2R11-S | Rabbit Anti-Porcine Circovirus 2 replicase protein (PCV2-rep/ORF1) antiserum |
| PCV2R15-R-10 | Recombinant (E.coli) Porcine Circovirus 2 replicase (PCV2-rep/ORF1) protein (his tag, 36 kDa) purified |
| AE-200250-1 | Recombivirus Q™ Porcine/Swine/Pig Anti-Circovirus 2 Capsid (PCV2-ORF2) IgG ELISA kit, Quantitative, 1x96 tests |
| AE-200250-5 | Recombivirus Q™ Porcine/Swine/Pig Anti-Circovirus 2 Capsid (PCV2-ORF2) IgG ELISA kit, Quantitative, 5x96 tests |
| AE-200260-1 | Recombivirus Q™ Porcine/Swine/Pig Anti-Porcine Circovirus 2 Replicase protein IgG ELISA kit (DIVA test), Quantitative, 1x96 tests |
| AE-200260-5 | Recombivirus Q™ Porcine/Swine/Pig Anti-Porcine Circovirus 2 Replicase protein IgG ELISA kit (DIVA test), Quantitative, 5x96 tests |

PCV2R11-S 141219P

Alpha Diagnostic Intl Inc., 6203 Woodlake Center Dr, San Antonio, TX 78244, USA; Email: service@4adi.com

India Contact:

Life Technologies (India) Pvt. Ltd.

306, Aggarwal City Mall, Opposite M2K Pitampura, Delhi - 110034 (INDIA). Ph: +91-11-42208000, 42208111, 42208222, Mobile: +91-9810521400, Fax: +91-11-42208444

Email: customerservice@lifetechindia.com Website: www.lifetechindia.com