

Product Data Sheet

Cat # RP-1026

Recombinant Human Stromal Cell-Derived Factor-1 beta (CXCL12)

Size: 10 ug

SDF-1 (stromal cell-derived factor-1) is small cytokine belonging to the chemokine family that is officially designated Chemokine (C-X-C motif) ligand 12 (CXCL12). It is produced in two forms, SDF-1 α /CXCL12a and SDF-1 β /CXCL12b, by alternate splicing of the same gene. Chemokines are characterized by the presence of four conserved cysteines, which form two disulfide bonds. The CXCL12 proteins belong to the group of CXC chemokines, whose initial pair of cysteines are separated by one intervening amino acid. CXCL12 is strongly chemotactic for lymphocytes and has been implicated as an important cell co-ordinator during development. During embryogenesis it directs the migration of hematopoietic cells from foetal liver to bone marrow. Mice which were knocked-out for CXCL12 gene were lethal before the birth or within just 1 hour of life. As another role, CXCL12a alters also the electrophysiology of neurons. CXCL12 was shown to be expressed in many tissues in mice (including brain, thymus, heart, lung, liver, kidney, spleen and bone marrow). The receptor for this chemokine is CXCR4, which was previously called fusin. This CXCL12-CXCR4 interaction used to be considered exclusive (unlike for other chemokines and their receptors), but recently it was suggested that CXCL12 is also bound by CXCR7 receptor. The gene for CXCL12 is located on human chromosome 10. In human and mouse both CXCL12 and CXCR4 show high identity of sequence: 99% and 90%, respectively.

Source: *Escherichia Coli*. Stromal Cell-Derived Factor-1 beta Human Recombinant produced in E.Coli is a non-glycosylated, Polypeptide chain containing 72 amino acids and having a molecular mass of 8508 Dalton. The SDF-1b is purified by proprietary chromatographic techniques. The CXCL12 protein was lyophilized from a concentrated (1mg/ml) sterile solution containing no additives. Protein quantization was carried out by two independent methods: 1. UV spectroscopy at 280 nm using the absorbency value of 1.06 as the extinction coefficient for a 0.1% (1mg/ml) solution. This value is calculated by the PC GENE computer analysis program of protein sequences (IntelliGenetics). 2. Analysis by RP-HPLC, using a standard solution of SDF-1 β as a Reference Standard.

Applications and Suggested Dilutions: Greater than 98.0% as determined by: (a) Analysis by RP-HPLC. (b) Analysis by SDS-PAGE. It is recommended to reconstitute the lyophilized Stromal Cell-Derived Factor-1b in sterile 18M Ω -cm H₂O not less than 100 μ g/ml, which can then be further diluted to other aqueous solutions. Users must optimize the appropriate concentration and conditions for each assay.

Storage and Stability: Lyophilized SDF-1b although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution CXCL12

should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). If supplied in powder then reconstitute it in 100 ul water for 1 mg/ml stock and store in liquid at 4oC for ~1 week or aliquots in suitable size and store at -20oC for long term storage. **Please prevent freeze-thaw cycles.**

Biological Activity: The specific activity as determined by its ability to chemo attracts human peripheral T cells activated with PHA and IL-2 using a concentration of 20-80 ng/ml.

Usage:

This item is for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals

RP-1027

120419A

Alpha Diagnostic Intl Inc., 6203 Woodlake Center Dr, S an Antonio, T X 7 8 24 4 , U S A;

India Contact:

Life Technologies (India) Pvt. Ltd.

306, Aggarwal City Mall, Opposite M2K Pitampura, Delhi – 110034 (INDIA). Ph: +91-11-42208000, 42208111, 42208222, Mobile: +91-9810521400 Fax: +91-11-42208444 Email: customerservice@lifetechindia.com Website: www.lifetechindia.com