

Product Data Sheet

Cat # RP-1037

Recombinant Mouse GRO/KC (CXCL1)

Size: 5 ug

Chemokine (C-X-C motif) ligand 1 (CXCL1) is a small cytokine belonging to the CXC chemokine family that was previously called GRO1 oncogene, Neutrophil-activating protein 3 (NAP-3) and melanoma growth stimulating activity, alpha (MSGA- α). It is secreted by human melanoma cells, has mitogenic properties and is implicated in melanoma pathogenesis. CXCL1 is expressed by macrophages, neutrophils and epithelial cells, and has neutrophil chemo attractant activity. CXCL1 plays a role in spinal cord development by inhibiting the migration of oligodendrocyte precursors and is involved in the processes of angiogenesis, inflammation, wound healing, and tumorigenesis. This chemokine elicits its effects by signaling through the chemokine receptor CXCR2. The gene for CXCL1 is located on human chromosome 4 amongst genes for other CXC chemokines.

Source: *Escherichia Coli*. KC Mouse Recombinant also known as N51 and GRO1 produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 72 amino acids and having a molecular mass of 7815 Dalton. The GRO-1 is purified by proprietary chromatographic techniques. The protein was lyophilized with no additives.

Applications and Suggested Dilutions: Greater than 95.0% as determined by: (a) Analysis by RP-HPLC. (b) Analysis by SDS-PAGE. It is recommended to reconstitute the lyophilized GRO1 Recombinant in sterile 18M Ω -cm H₂O not less than 100 μ g/ml, which can then be further diluted to other aqueous solutions. Users must optimize the appropriate concentration and conditions for each assay.

Storage and Stability: Lyophilized KC Mouse although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution CXCL1 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). If supplied in powder then reconstitute it in 100 ul water for 1 mg/ml stock and store in liquid at 4°C for ~1 week or aliquots in suitable size and store at -20°C for long term storage. **Please prevent freeze-thaw cycles.**

Biological Activity: The biological activity was determined by measuring the dose dependent mobilization of intracellular calcium (calcium flux) with human neutrophils. Significant calcium mobilization is observed with \geq 50ng/mL of recombinant mouse Gro1.

Protein content: Protein quantitation was carried out by two independent methods: 1. UV spectroscopy at 280 nm using the absorbency value of 0.03 as the extinction coefficient for a 0.1% (1mg/ml) solution. This value is

calculated by the PC GENE computer analysis program of protein sequences (IntelliGenetics). 2. Analysis by RP-HPLC, using a standard solution of KC as a Reference Standard

Usage:

This item is for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals

RP-1037 120419A

Alpha Diagnostic Intl Inc., 6203 Woodlake Center Dr, S an Antonio, T X 7 8 24 4 , U S A;

India Contact:

Life Technologies (India) Pvt. Ltd.

306, Aggarwal City Mall, Opposite M2K Pitampura, Delhi – 110034 (INDIA). Ph: +91-11-42208000, 42208111, 42208222, Mobile: +91-9810521400 Fax: +91-11-42208444 Email: customerservice@lifetechindia.com Website: www.lifetechindia.com