
Cat#	SP-100449-5
Description:	Pancreatic Polypeptide (31-36) (human) (AA: Leu-Thr-Arg-Pro-Arg-Tyr-NH ₂) (MW: 804)
Size:	5 mg
Purity:	>95%
Store:	Desiccated at -20oC.

Pancreatic hormone is synthesized in pancreatic islets of Langerhans and acts as a regulator of exocrine pancreatic secretion, biliary tract motility, and food intake. Its secretion in humans is increased after a protein meal, fasting, exercise, and acute [hypoglycemia](#) and is decreased by [somatostatin](#) and intravenous [glucose](#). Plasma PP has been shown to be reduced in conditions associated with increased food intake and elevated in [anorexia nervosa](#). It stimulates the gastric juice secretion, but inhibits the gastric secretion induced by pentagastrine. It is the antagonist of [cholecystokinin](#) and inhibits the pancreatic secretion which was stimulated by cholecystokinin. On fasting, PP seric concentration is 80 pg/ml; after the meal, it rises up from 8 to 10 times more; glucose and fats also induce PP's level increase, but on parenteral introduction of those substances, the level of hormones doesn't change. The administration of [atropine](#), the [vagotomy](#), blocks the PP's after-meal secretion. The excitation of the [vagus nerve](#), the administration of [gastrin](#), [secretin](#) or cholecystokinin induce PP secretion.

References:

Lonovics J., (1981). Arch Surg. 116 (10): 1256–64. Batterham, RL., (2003). The Journal of Clinical Endocrinology and Metabolism 88 (8): 3989–92. Asakawa, A., (1999). Peptides 20: 1445-1448.

Related products

- SP-100446-1 Pancreatic Polypeptide (bovine)
- SP-100447-1 Pancreatic Polypeptide (human]
- SP-100448-1 Pancreatic Polypeptide (rana temporaria)
- SP-100450-5 Pancreatic Polypeptide (31-36) (free acid) (human)
- SP-52291-1 Pancreatic Polypeptide, Rat;
- SP-52293-1 Pancreatic Polypeptide, Human;
- SP-55301-05 Pancreatic Polypeptide, Avian;

Rev. 150513SV