

Survivin WB Positive control

Cat. SURV11-C

Purified recombinant Human Survivin Protein WB +ve control

SIZE: 100 ul

The inhibitors of apoptosis proteins (IAPs) are a widely expressed gene family of apoptotic inhibitors. The central mechanisms of IAP apoptotic suppression appear to be through direct caspases and pro-caspase inhibition. Recently, a new human gene encoding a structurally and unique IAP designated **Survivin** has been identified. Survivin (human 142 aa, ~16.5 kDa, chromosome 17q25; Mouse TIAP/Survivin 140 aa) contains a single baculovirus IAP repeat and lacks a C-terminal RING finger. An isoform of **survivin, Beta-form**, that has a 23-aa insertion in the BIR-domain has been reported. Mouse survivin is also shown to be alternatively spliced into 3 forms: **Long 140-aa, 121-aa (missing the oiled-coil domain), and 40-aa form** (missing the IAP repeat and coiled domain). Survivin140 and 121-aa inhibits caspase-3 activity.

It has the property of oncofetal antigens: highly expressed in less-differentiated embryonic cells or rapidly dividing tumor cell but not in fully differentiated adult tissues. Elevated levels of Survivin are found in human fetal lung, liver, heart, kidney, and gastrointestinal tract. In mouse embryonic tissues, Survivin is detected in most tissues. High level of Survivin was found in most common human cancer, including cancers of the lung, colon, pancreas, prostate, and breast. Expression of Survivin also correlated with the presence of both p53 and bcl-2.

Source of Antigen and Antibodies

Human survivin was expressed as his-tagged fusion protein and purified. For **western blot +ve control (Cat # SURV11-C)**, it is supplied in SDS-PAGE sample buffer (reduced). Load ~10 ul/lane to visualize with appropriate antibodies (cat # SURV11-S or SURV11-A). Recombinant survivin is ~17 kDa. Store at -20oC in suitable aliquots. Avoid repeated thawing or heating.

General References: (1) Ambrosini G et al (1997) Nature Med. 3, 917-921; Altieri DC et al (1994) J Biol. Chem. 269, 3139-3142; Altieri DC et al (1994) Biochem. 33, 13848-13855; Adida C et al (1998) Am. J Pathol. 152, 43-49; Lu C-D et al (1998) Cancer Res. 58, 1808-1812; Li F et al (1998) Nature 396, 580-584; Kobayashi K et al (1998) Gene Accession # AB01389.

Published papers on ADI's Antibodies to Survivin (A complete and updated list posted at the web site)

Ikeguchi M, 2002, Diag. Mol. Pathol.11:33-40

Sasaki, 2002, Acta Neuropathol.

Guo , 2002, Blood 99, 3419-3429 **WB, IHC**

Shankar, 2001, J. Neurochem. 79, 426, msn cells/**WB**

Kanwar, 2001, J Natl Cancer Inst 93: 1541-1552,

Chantalat L 2000, Mol Cell, 6, 183-189, **WB,IF**

Okada E, 2001, Cancer Lett. 163, 109-116, **WB, IHC,IF**

Tran, J, 1999, BBRC 264, 781-788, **WB**

Storage

Short-term: unopened, undiluted vials for less than a week at 4oC.

Long-term: at -20C or below in suitable aliquots after reconstitution. Do not freeze and thaw and store working, diluted solutions.

Stability: 6-12 months at -20oC or below.

Shipping: 4oC for solutions and room temp for powder.

*This product is for *in vitro* research use only.

SURV11-C

80303A