™ *Phyto*Technology Laboratories, LLC™

Helping to Build a Better Tomorrow through Plant Science™

Product Information Sheet

L467

Linsmaier & Skoog Basal Medium w/ 30 g/L Sucrose and 7 g/L Agar

Synonym: Murashige & Skoog (MS) Medium with Minimal Organics (MSMO)

Properties

Form: Powder

Appearance: White to Yellow Powder Application: Plant Tissue Culture

Solubility: Water

Typical Working

41.43 g/L

Concentration:

Storage Temp: 2 – 6° C

Storage Temp of Preparation of concentrated solutions is not recommended as insoluble

Stock Solution: precipitates may form.

Other Notes: Contains the macro- and micronutrients and vitamins as described by

Linsmaier and Skoog (1965).

pH = 4.25 - 5.25

Formula (mg/L)

Ammonium Nitrate	1650
Boric Acid	6.2
Calcium Chloride, Anhydrous	332.2
Cobalt Chloride•6H ₂ O	0.025
Cupric Sulfate•5H ₂ O	0.025
Na ₂ EDTA•2H ₂ O	37.26
Ferrous Sulfate•7H ₂ O	27.8
Magnesium Sulfate, Anhydrous	180.7
Manganese Sulfate•H ₂ O	16.9
Molybdic Acid (Sodium Salt) •2H ₂ O	0.25

Potassium Iodide	0.83
Potassium Nitrate	1900
Potassium Phosphate, Monobasic	170
Zinc Sulfate•7H ₂ O	8.6
Agar	7000
myo-Inositol	100
Sucrose	30,000
Thiamine•HCl	0.4

Application Notes

Plant Tissue Culture Tested

Plant species: Originally tobacco; since then many other species

This medium is the standard Murashige & Skoog (MS) basal salts supplemented with Linsmaier and Skoog vitamins.

This medium is a subsequent optimization of the medium developed by Murashige and Skoog. Linsmaier's research on the optimization of vitamins first described by Murashige as essential. Linsmaier found that certain vitamins were not essential when Thiamine HCI was optimized

India Contact

Life Technologies (India) Pvt Ltd.



[™] *Phyto*Technology Laboratories, LLC™

Helping to Build a Better Tomorrow through Plant Science™

Your Molecular & Cell Technology Partner

Product Information Sheet

from 0.1 to 0.4 mg/L. Linsmaier also noted that Folic acid, p-Aminobenzoic Acid, L-Glutamine, and Ascorbic Acid all had a positive influence on the growth of Nicotiana callus but was not as essential as Thiamine•HCl and myo-Inositol.

References

Linsmaier, EM and F Skoog. 1965. Organic growth factor requirements of tobacco tissue culture. Physiol. Plant. 18:100-127.

Revised 2/2007

India Contact