


# **PhytoTechnology Laboratories**®

Helping to Build a Better Tomorrow through Plant Science™

Your Molecular & Cell Technology Partner

# **Product Information Sheet**



T8100 Triton X-100

Synonyms:Octoxynol; X-100; 4-(1,1,3,3-Tetramethylbutyl)phenyl-polyethylene glycol;<br/>*t*-Octylphenoxypolyethoxyethanol; Polyethylene glycol *tert*-octylphenyl ether<br/>CAS:9002-93-1<br/>Formula:C14H22O(C2H4O)n (n=9-10)Mol. Weight:Average 625

### **Properties**

Form: Liquid Appearance: Colorless to Light Yellow, Clear to Slightly Hazy Liquid Application: Molecular Biology Solubility: Miscible with Water Storage Temp: Room Temperature Typical Working Concentration: Varies

### **Application Notes**

Triton X-100 is a nonionic surfactant. It's often used to solubilize proteins in molecular biology applications.<sup>2, 3</sup> It has also been used to enhance the dispersal of compounds at hydrophobic-hydrophilic phase boundaries.

Please Note: It is the sole responsibility of the purchaser to determine the appropriateness of this product for the specific plants that are being cultured and applications that are being used.

### References

- 1. Merck **13**, 6793.
- Hearing, Vincent J., Walter G. Klingler, Thomas M. Ekel, and Paul M. Montague. 1976. Molecular weight estimation of Triton X-100 solubilized proteins by polyacrylamide gel electrophoresis. *Analytical Biochemistry*. 72(1-2):113-122.
- 3. Pappas, Peter W. 1982. Solubilization of the Membrane-Bound Enzymes of the Brush-Border Plasma Membrane of Hymenolepis diminuta (Cestoda) Using Nonionic Detergents. *Journal of Parasitology*. 68(4):588-592.

## **India Contact**

T8100-Info