

PhytoTechnology Laboratories®

Product Information Sheet

W898 White Basal Salt Mixture

Properties

Form:	Powder
Appearance:	White to Yellow
Application:	Plant Tissue Culture
Solubility:	Water
Typical Working	0.02 a/l
Concentration:	0.93 g/L
Storage Temp:	2 – 6°C
Storage Temp of	Preparation of concentrated solutions is not recommended as insoluble
Stock Solution:	precipitates may form.
Other Notes:	Contains the macro- and micronutrients as described by White (1963).

Formula

Boric Acid	1.5
Calcium Nitrate	208.5
Cupric Sulfate•5H ₂ O	0.001
Ferrous Sulfate	2.5
Magnesium Sulfate, Anhydrous	351.62
Manganese Sulfate•H ₂ O	5.31
Molybdenum Trioxide	0.0001

Potassium Chloride	65
Potassium Iodide	0.75
Potassium Nitrate	80
Sodium Phosphate Monobasic	16.5
Sodium Sulfate	200
Zinc Sulfate•7H ₂ O	3.0

Application Notes

Has been used to tissue culture: Sugarcane protoplasts (Thom & Maretzki, 1985) Brassica (Inomata, 1978) Carica papaya (Litz & Conover, 1981) Marchantia polymorpha (Ohta et al., 1977)

References

- Inomata, N. (1978). Production of interspecific hybrids in Brassica campestris x B. oleracea by culture in vitro of excised ovaries I. Development of excised ovaries in the crosses of various cultivars. 遺伝學雑誌, 53(3), 161-173.
- Litz, R. E., & Conover, R. A. (1981). In vitro polyembryony in Carica papaya L. ovules. Zeitschrift für Pflanzenphysiologie, 104(3), 285-288.

India Contact



Product Information Sheet

- Ohta, Y., Katoh, K., & Miyake, K. (1977). Establishment and growth characteristics of a cell suspension culture of Marchantia polymorpha L. with high chlorophyll content. Planta, 136(3), 229-232.
- Thom, M., & Maretzki, A. (1985). Evidence for a plasmalemma redox system in sugarcane. Plant physiology, 77(4), 873-876.
- White, PR. 1963. The Cultivation of Animal and Plant Cells, 2nd Edition, Ronald Press, NY, NY.

India Contact

Page 2 of 2 Revised 8/2015 CC